

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
9 August 2001 (09.08.2001)

PCT

(10) International Publication Number  
**WO 01/57070 A1**

(51) International Patent Classification<sup>7</sup>: **C07K 7/08**,  
7/52, A61K 38/00

[US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US).

(21) International Application Number: PCT/US01/03293

(74) Common Representative: **MERCK & CO., INC.**; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US).

(22) International Filing Date: 1 February 2001 (01.02.2001)

(25) Filing Language:

English

(81) Designated States (*national*): CA, JP, US.

(26) Publication Language:

English

(84) Designated States (*regional*): European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR).

(30) Priority Data:

60/179,967

3 February 2000 (03.02.2000)

US

(71) Applicant (*for all designated States except US*): **MERCK & CO., INC.** [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US).

Published:

— with international search report

(72) Inventor; and

(75) Inventor/Applicant (*for US only*): **BEDNAREK, Maria**

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*



**WO 01/57070 A1**

(54) Title: **MELANIN-CONCENTRATING HORMONE ANALOGS**

(57) Abstract: The present invention features truncated MCH analogs active at the MCH receptor. The truncated MCH analogs are optionally modified peptide derivatives of mammalian MCH. The analogs can bind to the MCH receptor and, preferably, bring about signal transduction. MCH analogs have a variety of different uses including being used as a research tool and being used therapeutically.

## TITLE OF THE INVENTION

## MELANIN-CONCENTRATING HORMONE ANALOGS

## CROSS-REFERENCE TO RELATED APPLICATIONS

- 5                   The present application claims priority to provisional application U.S. Serial No. 60/179,967, filed February 3, 2000, which is hereby incorporated by reference herein.

## BACKGROUND OF THE INVENTION

- 10                   Neuropeptides present in the hypothalamus play a major role in mediating the control of body weight. (Flier, *et al.*, 1998. *Cell*, 92, 437-440.) Melanin-concentrating hormone (MCH) produced in mammals is a cyclic 19-amino acid neuropeptide synthesized as part of a larger pre-prohormone precursor in the hypothalamus which also encodes neuropeptides NEI and NGE. (Nahon, *et al.*, 1990. 15 *Mol. Endocrinol.* 4, 632-637; Vaughan, *et al.*, U.S. Patent No. 5,049,655; and Vaughan, *et al.*, 1989. *Endocrinology* 125, 1660-1665.) MCH was first identified in salmon pituitary, and in fish MCH affects melanin aggregation thus affecting skin pigmentation. In trout and eels MCH has also been shown to be involved in stress induced or CRF-stimulated ACTH release. (Kawauchi, *et al.*, 1983. *Nature* 305, 20 321-323.)

- In humans two genes encoding MCH have been identified that are expressed in the brain. (Breton, *et al.*, 1993. *Mol. Brain Res.* 18, 297-310.) In mammals MCH has been localized primarily to neuronal cell bodies of the hypothalamus which are implicated in the control of food intake, including perikarya 25 of the lateral hypothalamus and zona incerta. (Knigge, *et al.*, 1996. *Peptides* 17, 1063-1073.)

- Pharmacological and genetic evidence suggest that the primary mode of MCH action is to promote feeding (orexigenic). MCH mRNA is up regulated in fasted mice and rats, in the *ob/ob* mouse and in mice with targeted disruption in the 30 gene for neuropeptide Y (NPY). (Qu, *et al.*, 1996. *Nature* 380, 243-247 and Erickson, *et al.*, 1996. *Nature* 381, 415-418.) Injection of MCH centrally (ICV) stimulates food intake and MCH antagonizes the hypophagic effects seen with  $\alpha$  melanocyte stimulating hormone ( $\alpha$ MSH). (Qu, *et al.*, 1996. *Nature* 380, 243-247.) MCH deficient mice are lean, hypophagic and have increased metabolic rate. 35 (Shimada, *et al.*, 1998. *Nature* 396, 670-673.) The administration of MCH has been

indicated to useful for promoting eating, appetite or the gain or maintenance of weight. (Maratos-Flier, U.S. Patent No. 5,849,708.)

MCH action is not limited to modulation of food intake as effects on the hypothalamic-pituitary-axis have been reported. (Nahon, 1994. *Critical Rev. in Neurobiol.* 8, 221-262.) MCH may be involved in the body response to stress as MCH can modulate the stress-induced release of CRF from the hypothalamus and ACTH from the pituitary. In addition, MCH neuronal systems may be involved in reproductive or maternal function.

## 10 SUMMARY OF THE INVENTION

The present invention features truncated MCH analogs active at the MCH receptor. The truncated MCH analogs are optionally modified peptide derivatives of mammalian MCH. The analogs can bind to the MCH receptor and, preferably, bring about signal transduction. MCH analogs have a variety of different uses including being used as a research tool and being used therapeutically.

Thus, a first aspect of the present invention describes a truncated MCH analog. The truncated MCH analog is an optionally modified peptide having the structure:

20 
$$Z1-X1-X2-X3-X4-X5-X6-X7-X8-X9-X10-X11-X12-X13-X14-X15-X16-X17-Z2$$

wherein X<sup>1</sup> is an optionally present amino acid that, if present, is either alanine, valine, leucine, isoleucine, proline, tryptophan, phenylalanine, methionine, glycine, serine, threonine, tyrosine, cysteine, asparagine, glutamine, lysine, arginine, histidine, aspartic acid, or glutamic acid, or a derivative thereof;

X<sup>2</sup> is an optionally present amino acid that, if present, is either alanine, valine, leucine, isoleucine, proline, tryptophan, phenylalanine, methionine, glycine, serine, threonine, tyrosine, cysteine, asparagine, glutamine, lysine, arginine, histidine, aspartic acid, or glutamic acid, or a derivative thereof;

X<sup>3</sup> is an optionally present amino acid that, if present, is either alanine, valine, leucine, isoleucine, proline, tryptophan, phenylalanine, methionine, glycine, serine, threonine, tyrosine, cysteine, asparagine, glutamine, lysine, arginine, histidine, aspartic acid or glutamic acid, or a derivative thereof;

X4 is an optionally present amino acid that, if present, is either alanine, valine, leucine, isoleucine, proline, tryptophan, phenylalanine, methionine, glycine, serine, threonine, tyrosine, cysteine, asparagine, glutamine, lysine, arginine, histidine, aspartic acid, glutamic acid, or norleucine, or a derivative thereof;

5 X5 is an optionally present amino acid that, if present, is either alanine, valine, leucine, isoleucine, proline, tryptophan, phenylalanine, methionine, glycine, serine, threonine, tyrosine, cysteine, asparagine, glutamine, lysine, arginine, histidine, aspartic acid or glutamic acid, or a derivative thereof;

10 X6 is an optionally present amino acid that, if present is either arginine, alanine, leucine, glycine, lysine, proline, asparagine, serine, histidine, nitroarginine, norleucine, or des-amino-arginine, or a derivative thereof,

X7 is either cysteine, homocysteine, or penicillamine, or a derivative thereof;

15 X8 is either methionine, norleucine, leucine, isoleucine, valine, methioninesulfoxide, or methioninesulfone, or a derivative thereof;

X9 is either leucine, isoleucine, valine, alanine, methionine, or 5-aminopentanoic acid, or a derivative thereof;

20 X10 is either glycine, alanine, leucine, norleucine, cyclohexylalanine, 5-aminopentanoic acid, asparagine, serine, sarcosine, isobutyric, or gamma-aminobutyric acid, or a derivative thereof;

X11 is either arginine, lysine, citrulline, histidine, or nitroarginine, or a derivative thereof;

X12 is either valine, leucine, isoleucine, alanine, or methionine, or a derivative thereof;

25 X13 is either phenylalanine, tyrosine, D-(*p*-benzoylphenylalanine), tryptophan, (1')- and (2')-naphthylalanine, cyclohexylalanine, or mono and multi-substituted phenylalanine wherein each substituent is independently selected from the group consisting of O-alkyl, alkyl, OH, NO<sub>2</sub>, NH<sub>2</sub>, F, I, and Br; or a derivative thereof;

30 X14 is either arginine, lysine, histidine, norarginine, or 5-aminopentanoic acid or a derivative thereof;

X15 is either proline, alanine, valine, leucine, isoleucine, methionine, sarcosine, or 5-aminopentanoic acid, or a derivative thereof;

X<sup>16</sup> is either cysteine, homocysteine, or penicillamine, or a derivative thereof;

X<sup>17</sup> is an optionally present amino acid that, if present, is either alanine, valine, leucine, isoleucine, proline, tryptophan, phenylalanine, methionine, glycine, serine, threonine, tyrosine, cysteine, asparagine, glutamine, lysine, arginine, histidine, aspartic acid or glutamic acid, or a derivative thereof;

Z<sup>1</sup> is an optionally present protecting group that, if present, is covalently joined to the N-terminal amino group;

Z<sup>2</sup> is an optionally present protecting group that, if present, is covalently joined to the C-terminal carboxy group;  
or a labeled derivative of said peptide;  
or a pharmaceutically acceptable salt of said peptide or of said labeled derivative.

Unless otherwise stated, those amino acids with a chiral center are provided in the L-enantiomer. Reference to "a derivative thereof" refers to the corresponding D-amino acid, N-alkyl-amino acid and  $\beta$ -amino acid.

Another aspect of the present invention describes a method of screening for a compound able to bind a MCH receptor. The method comprises the step of measuring the ability of the compound to effect binding of a truncated MCH analog to either the MCH receptor, a fragment of the receptor comprising a MCH binding site, a polypeptide comprising such a fragment, or a derivative of the polypeptide.

Another aspect of the present invention describes a method for increasing weight in a subject. The method comprises the step of administering to the subject an effective amount of a truncated MCH analog to produce a weight increase.

Another aspect of the present invention describes a method for increasing appetite in a subject. The method comprises the step of administering to the subject an effective amount of a truncated MCH analog to produce an appetite increase.

Another aspect of the present invention describes a method for measuring the ability of a compound to decrease weight or appetite in a subject. The method comprising the steps of:

a) administering to the subject an effective amount of a truncated MCH analog to produce a weight increase or appetite increase,

- b) administering the compound to the subject, and
- c) measuring the change in weight or appetite of the subject.

Other features and advantages of the present invention are apparent from the additional descriptions provided herein including the different examples.

- 5 The provided examples illustrate different components and methodology useful in practicing the present invention. The examples do not limit the claimed invention. Based on the present disclosure the skilled artisan can identify and employ other components and methodology useful for practicing the present invention.

## 10 BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 illustrates the results of an alanine scan where different amino acid residues of human MCH were replaced with alanine. The binding assay was performed by measuring inhibition of (<sup>125</sup>I-tyrosine, phenylalanine<sup>13</sup>)-MCH binding to cloned human MCH receptor (CHO clone). Cyclization sites (S-S) are indicated by

15 “\*”.

## DETAILED DESCRIPTION OF THE INVENTION

- Truncated MCH analogs contain about 10 to about 17 groups that are amino acids or amino acid derivatives. Using the present application as a guide truncated MCH analogs can be produced having significant MCH receptor activity, and in some cases having activity equal to or better than naturally occurring mammalian MCH. The smaller size of truncated MCH analogs offers advantages over longer-length MCH such as ease of synthesis and/or increased solubility in physiological buffers.

- 25 The MCH receptor is a G-protein coupled receptor that appears to be able to couple to Gi and Gq. Several references describe a receptor that is indicated to be a MCH receptor. (Chambers, *et al.*, 1999. *Nature* 400, 261-265; Saito, *et al.*, 1999. *Nature* 400, 265-269; Bächner, *et al.*, 1999. *FEBS Letters* 457:522-524; and Shimomura, *et al.*, 1999. *Biochemical and Biophysical Research Communications* 261, 622-626. These references are not admitted to be prior art to the claimed invention.)

The nucleic acid encoding for different variants of a MCH receptor is provided for by SEQ. ID. NOS. 1-3. The encoded amino acid sequences of the variants are provided by SEQ. ID. NOS. 4-6. The variants differ from each other by

the presence of additional amino acids at the N-terminal. One or more of these variants may be a physiological MCH receptor.

Significant MCH activity is preferably at least about 50%, at least about 75%, at least about 90%, or at least about 95%, the activity of mammalian MCH as determined by a binding assay or MCH receptor activity assay. Examples of such assays are provided below.

MCH analogs have a variety of different uses including being used as a research tool and being used therapeutically. Research tool applications generally involve the use of a truncated MCH analog and the presence of a MCH receptor or fragment thereof. The MCH receptor can be present in different environments such as a mammalian subject, a whole cell, and membrane fragments. Examples of research tool applications of truncated MCH analogs include screening for compounds active at the MCH receptor, determining the presence of the MCH receptor in a sample or preparation, examining the role or effect of MCH, and examining the role or effect of MCH antagonists.

Truncated MCH analogs can be used to screen for both MCH agonists and MCH antagonists. Screening for MCH agonists can be performed, for example, by using a truncated MCH analog in a competition experiment with test compounds. Screening for MCH antagonists can be performed, for example, by using a truncated MCH analog to produce MCH receptor activity and then measuring the ability of a compound to alter MCH receptor activity.

Truncated MCH analogs can be administered to a subject. A "subject" refers to a mammal including, for example, a human, a rat, a mouse, or a farm animal. Reference to subject does not necessarily indicate the presence of a disease or disorder. The term subject includes, for example, mammals being dosed with a truncated MCH analog as part of an experiment, mammals being treated to help alleviate a disease or disorder, and mammals being treated prophylactically to retard or prevent the onset of a disease or disorder.

MCH agonists can be used to achieve a beneficial effect in a subject. For example, a MCH agonist can be used to facilitate a weight gain, maintenance of weight and/or an appetite increase. Such effects are particularly useful for a patient having a disease or disorder, or under going a treatment, accompanied by weight loss. Examples of diseases or disorders accompanied by weight loss include anorexia, AIDS, wasting, cachexia, and frail elderly. Examples of treatments accompanied by weight loss include chemotherapy, radiation therapy, and dialysis.


MCH antagonists can also be used to achieve a beneficial effect in a patient. For example, a MCH antagonist can be used to facilitate weight loss, appetite decrease, weight maintenance, cancer (e.g., colon or breast) treatment, pain reduction, stress reduction and/or treatment of sexual dysfunction.

5

#### Truncated MCH Analogs

A truncated MCH analog is an optionally modified peptide having the structure:

10



wherein X<sup>1</sup> is an optionally present amino acid that, if present, is either alanine, valine, leucine, isoleucine, proline, tryptophan, phenylalanine, methionine, glycine, serine, threonine, tyrosine, cysteine, asparagine, glutamine, lysine, arginine, histidine, aspartic acid, or glutamic acid, or a derivative thereof; preferably, X<sup>1</sup> if present is aspartic acid or glutamic acid; more preferably, X<sup>1</sup> if present is aspartic acid; and more preferably, X<sup>1</sup> is not present;

X<sup>2</sup> is an optionally present amino acid that, if present, is either alanine, valine, leucine, isoleucine, proline, tryptophan, phenylalanine, methionine, glycine, serine, threonine, tyrosine, cysteine, asparagine, glutamine, lysine, arginine, histidine, aspartic acid, or glutamic acid, or a derivative thereof; preferably, X<sup>2</sup> if present is phenylalanine or tyrosine; more preferably, X<sup>2</sup> if present is phenylalanine; and more preferably, X<sup>2</sup> is not present;

X<sup>3</sup> is an optionally present amino acid that, if present, is either alanine, valine, leucine, isoleucine, proline, tryptophan, phenylalanine, methionine, glycine, serine, threonine, tyrosine, cysteine, asparagine, glutamine, lysine, arginine, histidine, aspartic acid or glutamic acid, or a derivative thereof; preferably, X<sup>3</sup> if present is aspartic acid or glutamic acid; more preferably, X<sup>3</sup> if present is aspartic acid; and more preferably, X<sup>3</sup> is not present;

X<sup>4</sup> is an optionally present amino acid that, if present, is either alanine, valine, leucine, isoleucine, proline, tryptophan, phenylalanine, methionine, glycine, serine, threonine, tyrosine, cysteine, asparagine, glutamine, lysine, arginine, histidine, aspartic acid, glutamic acid, or norleucine, or a derivative thereof; preferably, X<sup>4</sup> if

present is methionine, leucine, isoleucine, valine, alanine or norleucine; more preferably, X<sup>4</sup> if present is methionine; and more preferably, X<sup>4</sup> is not present;

X<sup>5</sup> is an optionally present amino acid that, if present, is either alanine, valine, leucine, isoleucine, proline, tryptophan, phenylalanine, methionine, glycine, serine, threonine, tyrosine, cysteine, asparagine, glutamine, lysine, arginine, histidine, aspartic acid or glutamic acid, or a derivative thereof; preferably, X<sup>5</sup> if present is leucine, methionine, isoleucine, valine or alanine; more preferably, X<sup>5</sup> if present is leucine; and more preferably, X<sup>5</sup> is not present;

X<sup>6</sup> is an optionally present amino acid that, if present is either arginine, alanine, leucine, glycine, lysine, proline, asparagine, serine, histidine, nitroarginine, norleucine, or des-amino-arginine, or a derivative thereof; preferably X<sup>6</sup> is not present or is either arginine, D-arginine, D-norleucine, D-proline, D-serine, or D-asparagine; more preferably X<sup>6</sup> is arginine or D-arginine;

X<sup>7</sup> is either cysteine, homocysteine, or penicillamine, or a derivative thereof; preferably, X<sup>7</sup> is cysteine;

X<sup>8</sup> is either methionine, norleucine, leucine, isoleucine, valine, methioninesulfoxide, or methioninesulfone, or a derivative thereof; preferably, X<sup>8</sup> is methionine, norleucine, or N-methyl norleucine;

X<sup>9</sup> is either leucine, isoleucine, valine, alanine, methionine, or 5-aminopentanoic acid, or a derivative thereof; preferably, X<sup>9</sup> is leucine;

X<sup>10</sup> is either glycine, alanine, leucine, norleucine, cyclohexylalanine, 5-aminopentanoic acid, gamma-aminobutyric acid, asparagine, serine, sarcosine, or isobutyric or a derivative thereof; preferably, X<sup>10</sup> is either glycine, alanine, leucine, norleucine, asparagine, serine, D-norleucine, D-proline, gamma-aminobutyric acid, or sarcosine; more preferably X<sup>10</sup>, is either glycine, leucine, norleucine, asparagine, or serine;

X<sup>11</sup> is either arginine, lysine, citrulline, histidine, or nitroarginine, or a derivative thereof; preferably, X<sup>11</sup> is arginine;

X<sup>12</sup> is either valine, leucine, isoleucine, alanine, or methionine, or a derivative thereof; preferably, X<sup>12</sup> is valine;

X<sup>13</sup> is either phenylalanine, tyrosine, D-(*p*-benzoylphenylalanine), tryptophan, (1')- and (2')-naphthylalanine, cyclohexylalanine, or mono and multi-substituted phenylalanine wherein each substituent is independently selected from the group consisting of O-alkyl, alkyl, OH, NO<sub>2</sub>, NH<sub>2</sub>, F, I, and Br; or a derivative

thereof; preferably, X<sup>13</sup> is phenylalanine, (2')naphthylalanine, p-fluoro-phenylalanine, tyrosine, or cyclohexylalanine;

X<sup>14</sup> is either arginine, lysine, histidine or norarginine, or 5-aminopentanoic acid, or a derivative thereof; preferably, X<sup>14</sup> is arginine;

5 X<sup>15</sup> is either proline, alanine, valine, leucine, isoleucine, methionine, sarcosine, or 5-aminopentanoic acid, or a derivative thereof; preferably, X<sup>15</sup> is proline or sarcosine;

X<sup>16</sup> is either cysteine, homocysteine, or penicillamine, or a derivative thereof; preferably, X<sup>16</sup> is cysteine or D-cysteine;

10 X<sup>17</sup> is an optionally present amino acid that, if present, is either alanine, valine, leucine, isoleucine, proline, tryptophan, phenylalanine, methionine, glycine, serine, threonine, tyrosine, cysteine, asparagine, glutamine, lysine, arginine, histidine, aspartic acid or glutamic acid, or a derivative thereof; preferably, X<sup>17</sup> if present is tyrosine or tryptophan; more preferably X<sup>17</sup> is not present;

15 Z<sup>1</sup> is an optionally present protecting group that, if present, is covalently joined to the N-terminal amino group;

Z<sup>2</sup> is an optionally present protecting group that, if present, is covalently joined to the C-terminal carboxy group;

or a labeled derivative of said peptide;

20 or a pharmaceutically acceptable salt of said peptide or of said labeled derivative.

The present invention is meant to comprehend diastereomers as well as their racemic and resolved enantiomerically pure forms. Truncated MCH analogs can contain D-amino acids, L-amino acids or a combination thereof. Preferably, amino  
25 acids present in a truncated MCH analog are the L-enantiomer.

In different embodiments, MCH analogs contain a preferred (or more preferred) group at one or more different locations. More preferred embodiments contain preferred (or more preferred) groups in each of the different locations.

A protecting group covalently joined to the N-terminal amino group  
30 reduces the reactivity of the amino terminus under *in vivo* conditions. Amino protecting groups include optionally substituted -C<sub>1-10</sub> alkyl, optionally substituted -C<sub>2-10</sub> alkenyl, optionally substituted aryl, -C<sub>1-6</sub> alkyl optionally substituted aryl, -C(O)-(CH<sub>2</sub>)<sub>1-6</sub>-COOH, -C(O)-C<sub>1-6</sub> alkyl, -C(O)-optionally substituted aryl,

-C(O)-O-C<sub>1-6</sub> alkyl, or -C(O)-O-optionally substituted aryl. Preferably, the amino terminus protecting group is acetyl, propyl, succinyl, benzyl, benzyloxycarbonyl or *t*-butyloxycarbonyl.

5 A protecting group covalently joined to the C-terminal carboxy group reduces the reactivity of the carboxy terminus under *in vivo* conditions. The carboxy terminus protecting group is preferably attached to the  $\alpha$ -carbonyl group of the last amino acid. Carboxy terminus protecting groups include amide, methylamide, and ethylamide.

"Alkyl" refers to carbon atoms joined by carbon-carbon single bonds.

10 The alkyl hydrocarbon group may be straight-chain or contain one or more branches or cyclic groups. Preferably, the alkyl group is 1 to 4 carbons in length. Examples of alkyl include methyl, ethyl, propyl, isopropyl, cyclopropyl, butyl, and *t*-butyl. Alkyl substituents are selected from the group consisting of halogen (preferably -F or -Cl) -OH, -CN, -SH, -NH<sub>2</sub>, -NO<sub>2</sub>, -C<sub>1-2</sub> alkyl substituted with 1 to 6 halogens (preferably  
15 -F or -Cl, more preferably -F), -CF<sub>3</sub>, -OCH<sub>3</sub>, or -OCF<sub>3</sub>.

"Alkenyl" refers to a hydrocarbon group containing one or more carbon-carbon double bonds. The alkenyl hydrocarbon group may be straight-chain or contain one or more branches or cyclic groups. Preferably, the alkenyl group is 2 to 4 carbons in length. Alkenyl substituents are selected from the group consisting of  
20 halogen (preferably -F or -Cl), -OH, -CN, -SH, -NH<sub>2</sub>, -NO<sub>2</sub>, -C<sub>1-2</sub> alkyl substituted with 1 to 5 halogens (preferably -F or -Cl, more preferably -F), -CF<sub>3</sub>, -OCH<sub>3</sub>, or -OCF<sub>3</sub>.


"Aryl" refers to an optionally substituted aromatic group with at least one ring having a conjugated pi- electron system, containing up to two  
25 conjugated or fused ring systems. Aryl includes carbocyclic aryl, heterocyclic aryl and biaryl groups. Preferably, the aryl is a 5 or 6 membered ring, more preferably benzyl. Aryl substituents are selected from the group consisting of -C<sub>1-4</sub> alkyl, -C<sub>1-4</sub> alkoxy, halogen (preferably -F or -Cl), -OH, -CN, -SH, -NH<sub>2</sub>, -NO<sub>2</sub>, -C<sub>1-2</sub> alkyl substituted with 1 to 5 halogens (preferably -F or -Cl, more preferably -F), -CF<sub>3</sub>, or  
30 -OCF<sub>3</sub>.

A labeled derivative indicates the alteration of a substituent with a detectable label. Examples of detectable labels include luminescent, enzymatic, and radioactive labels. A preferred radiolabel is <sup>125</sup>I. Both the type of label and the position of the label can effect MCH activity. Labels should be selected so as not to  
35 substantially alter the activity of the truncated MCH analog at the MCH receptor. The

effect of a particular label on MCH activity can be determined using assays measuring MCH activity and/or binding.

In naturally occurring full length MCH, alteration of the tyrosine at position 13 by labeling with  $^{125}\text{I}$  substantially effects MCH activity. (Drozd, *et al.*, 1995. *FEBS letters* 359, 199-202.)  $^{125}\text{I}$  labeled analogs of full length mammalian MCH having substantial activity can be produced, for example, by replacing the tyrosine at position 13 with a different group, then replacing valine at position 19 with tyrosine, and labeling the tyrosine. Examples of such analogs include [ $^{125}\text{I}$ ][Phe<sup>13</sup>, Try<sup>19</sup>]-MCH and (D-(*p*-benzoylphenylalanine)<sup>13</sup>, tyrosine<sup>19</sup>)-MCH. (Drozd, *et al.*, 1995. *FEBS letters* 359, 199-202, 1995; and Drozd, *et al.*, *J. Peptide Sci.* 5, 234-242, 1999.)

In preferred embodiments the optionally modified peptide has the structure:



wherein the different groups, and preferred groups, are as described above.

In different embodiments the truncated MCH analog is a peptide of SEQ. ID. NOS. 7, 8, 9, or 10, a labeled derivative of said peptide or a pharmaceutically acceptable salt of said peptide or of said labeled derivative. SEQ. ID. NOS. 7-12 are made up of L-amino acids and have the following sequences ("\*" indicates cyclization (S-S)):

- SEQ. ID. NO. 7: Ac-Arg-Cys-Met-Leu-Gly-Arg-Val-Tyr-Arg-Pro-Cys-amide;
- SEQ. ID. NO. 8: Ac-Arg-Cys-Met-Leu-Gly-Arg-Val-Phe-Arg-Pro-Cys-Tyr-amide;
- SEQ. ID. NO. 9: Ac-Cys-Met-Leu-Gly-Arg-Val-Tyr-Arg-Pro-Cys-amide;
- SEQ. ID. NO. 10: Asp-Phe-Asp-Met-Leu-Arg-Cys-Met-Leu-Gly-Arg-Val-Tyr-Arg-Pro-Cys-amide;

SEQ. ID. NO. 12: Ac-Cys-Met-Leu-Gly-Arg-Val-Tyr-Arg-Pro-Cys-Trp-Gln-Val;

SEQ. ID. NO. 13:

Asp-Phe-Asp-Nle-Leu-Arg-Cys-Nle-Leu-Gly-Arg-Val-Tyr-Arg-Pro-Cys-Trp-Gln-Val;

SEQ. ID. NO. 14:

Asp-Phe-Ala-Met-Leu-Arg-Cys-Met-Leu-Gly-Arg-Val-Phe-Arg-Pro-Cys-Trp-Gln-Tyr.

In additional embodiments the peptide has a sequence selected from the group consisting of SEQ. ID. NOs. 7, 8, 10, 15, 24, 25, 27, 28, 30-49, 51, 52, 56, 57, 61, 62, 63, 65-67, 69-72, and 77, is a labeled derivative of said peptide or a pharmaceutically acceptable salt of said peptide or of said labeled derivative. Preferred sequences are those with an IC<sub>50</sub> less than 0.3 nM, preferably less than 0.1 nM; and/or those having a % activation greater than about 90%, preferably greater than 100%. Examples of preferred sequences are provided in Example 4, Tables 1-7.

Truncated MCH analogs can be produced using techniques well known in the art. For example, a polypeptide region of a truncated MCH analog can be chemically or biochemically synthesized and, if desired modified to produce a blocked N-terminus and/or blocked C-terminus. Techniques for chemical synthesis of polypeptides are well known in the art. (See *e.g.*, Vincent, in *Peptide and Protein Drug Delivery*, New York, N.Y., Dekker, 1990.) Examples of techniques for biochemical synthesis involving the introduction of a nucleic acid into a cell and expression of nucleic acids are provided in Ausubel, *Current Protocols in Molecular Biology*, John Wiley, 1987-1998, and Sambrook, *et al.*, in *Molecular Cloning, A Laboratory Manual*, 2<sup>nd</sup> Edition, Cold Spring Harbor Laboratory Press, 1989.

#### MCH Receptor Binding Assay

Assays measuring the ability of a compound to bind a MCH receptor employ a MCH receptor, a fragment of the receptor comprising a MCH binding site, a

polypeptide comprising such a fragment, or a derivative of the polypeptide.  
Preferably, the assay uses the MCH receptor or a fragment thereof.

5 A polypeptide comprising a MCH receptor fragment that binds MCH can also contain one or more polypeptide regions not found in a MCH receptor. A derivative of such a polypeptide comprises a MCH receptor fragment that binds MCH along with one or more non-peptide components.

10 The MCH receptor amino acid sequence involved in MCH binding can be readily identified using labeled MCH or truncated MCH analogs and different receptor fragments. Different strategies can be employed to select fragments to be tested to narrow down the binding region. Examples of such strategies include testing consecutive fragments about 15 amino acids in length starting at the N-terminus, and testing longer length fragments. If longer length fragments are tested, a fragment binding MCH can be subdivided to further locate the MCH binding region. Fragments used for binding studies can be generated using recombinant nucleic acid techniques.

15 Binding assays can be performed using individual compounds or preparations containing different numbers of compounds. A preparation containing different numbers of compounds having the ability to bind to the MCH receptor can be divided into smaller groups of compounds that can be tested to identify the compound(s) binding to the MCH receptor. In an embodiment of the present invention a test preparation containing at least 10 compounds is used in a binding assay.

20 Binding assays can be performed using recombinantly produced MCH receptor polypeptides present in different environments. Such environments include, for example, cell extracts and purified cell extracts containing the MCH receptor polypeptide expressed from recombinant nucleic acid or naturally occurring nucleic acid; and also include, for example, the use of a purified MCH receptor polypeptide produced by recombinant means or from naturally occurring nucleic acid which is introduced into a different environment.

30

#### Screening for MCH Receptor Active Compounds

Screening for MCH active compounds is facilitated using a recombinantly expressed MCH receptor. Using recombinantly expressed MCH receptor polypeptides offers several advantages such as the ability to express the receptor in a defined cell system so that response to MCH receptor active compounds

35

can more readily be differentiated from responses to other receptors. For example, the MCH receptor can be expressed in a cell line such as HEK 293, COS 7, and CHO not normally expressing the receptor by an expression vector, wherein the same cell line without the expression vector can act as a control.

5                Screening for MCH receptor active compounds is facilitated through the use of a truncated MCH analog in the assay. The use of a truncated MCH analog in a screening assay provides for MCH receptor activity. The effect of test compounds on such activity can be measured to identify, for example, allosteric modulators and antagonists. Additionally, such assays can be used to identify  
10 agonists.

                 MCH receptor activity can be measured using different techniques such as detecting a change in the intracellular conformation of the MCH receptor, Gi or Gq activity, and/or intracellular messengers. Gi activity can be measured using techniques well known in the art such as a melonaphore assay, assays measuring  
15 cAMP production, inhibition of cAMP accumulation, and binding of  $^{35}\text{S}$ -GTP. cAMP can be measured using different techniques such as radioimmunoassay and indirectly by cAMP responsive gene reporter proteins.

                 Gq activity can be measured using techniques such as those measuring intracellular  $\text{Ca}^{2+}$ . Examples of techniques well known in the art that can be  
20 employed to measure  $\text{Ca}^{2+}$  include the use of dyes such as Fura-2 and the use of  $\text{Ca}^{2+}$ -bioluminescent sensitive reporter proteins such as aequorin. An example of a cell line employing aequorin to measure G-protein activity is HEK293/aeq17. (Button, *et al.*, 1993. *Cell Calcium* 14, 663-671, and Feighner, *et al.*, 1999. *Science* 284, 2184-2188, both of which are hereby incorporated by reference herein.)

25                Chimeric receptors containing a MCH binding region functionally coupled to a G protein can also be used to measure MCH receptor activity. A chimeric MCH receptor contains an N-terminal extracellular domain; a transmembrane domain made up of transmembrane regions, extracellular loop regions, and intracellular loop regions; and an intracellular carboxy terminus.  
30 Techniques for producing chimeric receptors and measuring G protein coupled responses are provided for in, for example, International Application Number WO 97/05252, and U.S. Patent Number 5,264,565, both of which are hereby incorporated by reference herein.

### Weight or Appetite Alteration

Truncated MCH analogs can be used in methods to increase or maintain weight and/or appetite in a subject. Such methods can be used, for example, as part of an experimental protocol examining the effects of MCH antagonists, to achieve a beneficial effect in a subject and/or to further examine the physiological effects of MCH.

Experimental protocols examining the effects of MCH antagonists can be performed, for example, by using a sufficient amount of a truncated MCH analog to produce a weight or appetite increase in a subject and then examining the effect of a test compound. Changes in weight and appetite can be measured using techniques well known in the art.

Increasing weight or appetite can be useful for maintaining weight or producing a weight or appetite gain in an under weight subject, or in a patient having a disease or undergoing treatment that effects weight or appetite. In addition, for example, farm animals such as pigs, cows and chickens can be treated to gain weight.

Under weight subjects include those having a body weight about 10% or less, 20% or less, or 30% or less, than the lower end of a "normal" weight range or Body Mass Index ("BMI"). "Normal" weight ranges are well known in the art and take into account factors such as a patient age, height, and body type.

BMI measures your height/weight ratio. It is determined by calculating weight in kilograms divided by the square of height in meters. The BMI "normal" range is 19-22.

### Administration

Truncated MCH analogs can be formulated and administered to a subject using the guidance provided herein along with techniques well known in the art. The preferred route of administration ensures that an effective amount of compound reaches the target. Guidelines for pharmaceutical administration in general are provided in, for example, *Remington's Pharmaceutical Sciences 18<sup>th</sup> Edition*, Ed. Gennaro, Mack Publishing, 1990, and *Modern Pharmaceutics 2<sup>nd</sup> Edition*, Eds. Banker and Rhodes, Marcel Dekker, Inc., 1990, both of which are hereby incorporated by reference herein.

Truncated MCH analogs can be prepared as acidic or basic salts. Pharmaceutically acceptable salts (in the form of water- or oil-soluble or dispersible products) include conventional non-toxic salts or the quaternary ammonium salts that

are formed, *e.g.*, from inorganic or organic acids or bases. Examples of such salts include acid addition salts such as acetate, adipate, alginate, aspartate, benzoate, benzenesulfonate, bisulfate, butyrate, citrate, camphorate, camphorsulfonate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, fumarate, glucoheptanoate, glycerophosphate, hemisulfate, heptanoate, hexanoate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxyethanesulfonate, lactate, maleate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, oxalate, pamoate, pectinate, persulfate, 3-phenylpropionate, picrate, pivalate, propionate, succinate, tartrate, thiocyanate, tosylate, and undecanoate; and base salts such as ammonium salts, alkali metal salts such as sodium and potassium salts, alkaline earth metal salts such as calcium and magnesium salts, salts with organic bases such as dicyclohexylamine salts, N-methyl-D-glucamine, and salts with amino acids such as arginine and lysine.

Truncated MCH analogs can be administered using different routes including oral, nasal, by injection, transdermal, and transmucosally. Active ingredients to be administered orally as a suspension can be prepared according to techniques well known in the art of pharmaceutical formulation and may contain microcrystalline cellulose for imparting bulk, alginic acid or sodium alginate as a suspending agent, methylcellulose as a viscosity enhancer, and sweeteners/flavoring agents. As immediate release tablets, these compositions may contain microcrystalline cellulose, dicalcium phosphate, starch, magnesium stearate and lactose and/or other excipients, binders, extenders, disintegrants, diluents and lubricants.

Truncated MCH analogs may also be administered in intravenous (both bolus and infusion), intraperitoneal, subcutaneous, topical with or without occlusion, or intramuscular form. When administered by injection, the injectable solution or suspension may be formulated using suitable non-toxic, parenterally-acceptable diluents or solvents, such as Ringer's solution or isotonic sodium chloride solution, or suitable dispersing or wetting and suspending agents, such as sterile, bland, fixed oils, including synthetic mono- or diglycerides, and fatty acids; including oleic acid.

Suitable dosing regimens are preferably determined taking into factors well known in the art including type of subject being dosed; age, weight, sex and medical condition of the subject; the route of administration; the renal and hepatic function of the subject; the desired effect; and the particular compound employed.

Optimal precision in achieving concentrations of drug within the range that yields efficacy without toxicity requires a regimen based on the kinetics of the drug's availability to target sites. This involves a consideration of the distribution, equilibrium, and elimination of a drug. The daily dose for a subject is expected to be  
5 between 0.01 and 1,000 mg per subject per day.

Truncated MCH analogs can be provided in kit. Such a kit typically contains an active compound in dosage forms for administration. A dosage form contains a sufficient amount of active compound such that a weight or appetite increase can be obtained when administered to a subject during regular intervals, such  
10 as 1 to 6 times a day, during the course of 1 or more days. Preferably, a kit contains instructions indicating the use of the dosage form for weight or appetite increase and the amount of dosage form to be taken over a specified time period.

### EXAMPLES

15 Examples are provided below to further illustrate different features of the present invention. The examples also illustrate useful methodology for practicing the invention. These examples do not limit the claimed invention.

#### Example 1: Synthesis of MCH Analogs

20 MCH analogs were produced using the procedures described below and varying the stepwise addition of amino acid groups. Other procedures for producing and modifying peptides are well known in the art.

Elongation of peptidyl chains on 4-(2',4'-dimethoxyphenyl)-Fmoc-aminomethyl)-phenoxy resin and the acetylation of the N-terminal amino groups of  
25 the peptides was performed on a 431A ABI peptide synthesizer. Manufacture-supplied protocols were applied for coupling of the hydroxybenzotriazole esters of amino acids in N-methylpyrrolidone (NMP). The fluorenylmethyloxycarbonyl (Fmoc) group was used as a semipermanent alpha-amino protecting group, whereas the side chains protecting groups were: *tert*-butyl for aspartic acid and tyrosine,  
30 2,2,4,6,7-pentamethyldihydrobenzofuran-5-sulfonyl (Pbf) for arginine, and trityl for cysteine.

Peptides were cleaved from the resin with TFA containing 5 % of anisole. After 2 hours at room temperature the resin was filtered, washed with TFA and the combined filtrates were evaporated to dryness in vacuo. The residue was

triturerated with ether, the precipitate which formed was filtered off. washed with ether, and dried.

Crude peptides were dissolved in 5 % acetic acid in water, and the pH of the solutions were adjusted to ca. 8.2 with diluted ammonium hydroxide. The reaction mixtures were stirred vigorously while 0.05 % solution of potassium ferricyanide ( $K_3Fe(CN)_6$ ) in water was added dropwise till the reaction mixture remained yellow for about 5 minutes. After an additional 20 minutes oxidation was terminated with ca. 1 ml of acetic acid and the reaction mixtures were lyophilized.

Crude lyophilized peptides were analyzed by analytical reverse-phase high-pressure liquid chromatography (RP HPLC) on a C18 Vydac column attached to a Waters 600E system with automatic Wisp 712 injector and 991 Photodiode Array detector. A standard gradient system of 0-100% buffer B in 30 minutes was used for analysis: buffer A was 0.1% trifluoroacetic acid in water and buffer B was 0.1% trifluoroacetic acid in acetonitrile. HPLC profiles were recorded at 210 nm and 280 nm. Preparative separations were performed on a Waters Delta Prep 4000 system with a semipreparative C18 RP Waters column. The above-described solvent system of water and acetonitrile, in a gradient of 20-80 % buffer B in 60 minutes, was used for separation. The chromatographically homogenous compounds were analyzed by electrospray mass spectrometry.

20

#### Example 2: Aequorin Bioluminescence Functional Assay

The aequorin bioluminescence assay is a reliable test for measuring the activity of G protein-coupled receptors that couple through the  $G\alpha$  protein subunit family consisting of  $Gq$  and  $G11$  and leads to the activation of phospholipase C, mobilization of intracellular calcium and activation of protein kinase C.

Measurement of MCH receptor activity in the aequorin-expressing stable reporter cell line 293-AEQ17 (Button *et al.*, *Cell Calcium* 14:663-671, 1993) was performed using a Luminoskan RT luminometer (Labsystems Inc., Gaithersburg, MD). 293-AEQ17 cells ( $8 \times 10^5$  cells plated 18 hours before transfection in a T75 flask) were transfected with 22  $\mu$ g of human MCH receptor plasmid using 264  $\mu$ g lipofectamine. The open reading frame cDNA (SEQ. ID. NO. 1) encoding the human MCH receptor inserted in the mammalian expression vector pcDNA-3 (Invitrogen, Carlsbad, CA) was used for expression studies. Following approximately 40 hours of expression the apo-aequorin in the cells was charged for 4 hours with coelenterazine (10  $\mu$ M) under reducing conditions (300  $\mu$ M reduced glutathione) in ECB buffer (140

35

mM NaCl, 20 mM KCl, 20 mM HEPES-NaOH [pH=7.4], 5 mM glucose, 1 mM MgCl<sub>2</sub>, 1 mM CaCl<sub>2</sub>, 0.1 mg/ml bovine serum albumin).

The cells were harvested, washed once in ECB medium and resuspended to 500,000 cells/ml. 100 µl of cell suspension (corresponding to 5x10<sup>4</sup> cells) was then injected into the test plate containing MCH or MCH analogs, and the integrated light emission was recorded over 30 seconds, in 0.5 second units. 20 µL of lysis buffer (0.1% final Triton X-100 concentration) was then injected and the integrated light emission recorded over 10 seconds, in 0.5 second units. The "fractional response" values for each well were calculated by taking the ratio of the integrated response to the initial challenge to the total integrated luminescence including the Triton X-100 lysis response.

### Example 3: Radiolabeled MCH-R Binding Assay

Activity of truncated MCH analogs was assayed by measuring the ability of the analog to inhibit binding of [<sup>125</sup>I]-human MCH (Phe<sup>13</sup>, Tyr<sup>19</sup> substituted) to membranes prepared from cells stably expressing the human MCH receptor. Human MCH (Phe<sup>13</sup>, Tyr<sup>19</sup> substituted) used in the assay was radiolabeled with <sup>125</sup>I at <sup>19</sup>Tyr to a specific activity of ~2000 Ci/mmol (NEN Life Science Products, Boston, MA).

Cell membranes were prepared on ice. Each T-75 flask was rinsed twice with 10 ml of Enzyme-free Cell Dissociation Buffer (Specialty Media, Lavallete, NJ), and the cell monolayer was detached in an additional 10 ml of Enzyme-free Cell Dissociation Buffer by incubation at room temperature for 10 minutes. Dissociated cells were centrifuged (500 x g for 10 minutes at 4°C), resuspended in 5 ml homogenization buffer (10 mM Tris-HCl, pH 7.4, 0.01 mM Pefabloc, 10 µM phosphoramidon, 40 µg/ml bacitracin) and then homogenized using a glass homogenizer (10-15 strokes). The homogenate was centrifuged for 10 minutes (1,000 x g at 4°C). The resulting supernatant was then centrifuged at 38,700 x g for 15 minutes at 4°C. Pelleted membranes were resuspended (passed through 25 gauge needle 5 times), snap-frozen on liquid nitrogen, and stored at -80°C until use.

Binding was performed in a 96-well filter assay or Scintillation Proximity Assay (SPA)-based format using cell membranes from a stable CHO or HEK-293 cell line expressing the MCH receptor. For the filter assay, reactions were performed at 20°C for 1 hour in a total volume of 0.2 ml containing: 0.05 ml of membrane suspension (~3 µg protein), 0.02 ml of [<sup>125</sup>I]-human MCH (Phe<sup>13</sup>, Tyr<sup>19</sup>

substituted; 30 pM), 0.01 ml of competitor and 0.12 ml of binding buffer (50 mM Tris-HCl, pH 7.4, 10 mM MgCl<sub>2</sub>, 2 mM EDTA, 200 µg/ml bacitracin, 1 µM phosphoramidon).

Bound radioligand was separated by rapid vacuum filtration (Packard  
5 Filtermate 96-well cell harvester) through GF/C filters pretreated for 1 hour with 1 % polyethylenimine. After application of the membrane suspension to the filter, the filters were washed 3 times with 3 ml each of ice-cold 50 mM Tris-HCl, pH 7.4, 10 mM MgCl<sub>2</sub>, 2 mM EDTA, 0.04 % Tween 20 and the bound radioactivity on the filters was quantitated by scintillation counting (TopCount device). Specific binding  
10 (>80 % of total) is defined as the difference between total binding and non-specific binding conducted in the presence of 100 nM unlabeled human MCH.

For the SPA-based assay, WGA-PVT beads (NEN Life Sciences Products) were resuspended in Dulbecco's PBS with calcium and magnesium (500 mg beads in 4 ml PBS). For each 96-well assay plate, 0.18 ml of beads was pre-  
15 coated with MCH receptor by mixing with 0.2 ml MCH receptor CHO cell membranes (~ 0.2-4 mg protein) and 1.5 ml SPA assay buffer (50 mM Tris-HCl, pH 7.4, 10 mM MgCl<sub>2</sub>, 2 mM EDTA, 0.1 % BSA, 12 % glycerol). The suspension was mixed gently for 20 minutes, 12.3 ml of assay buffer and protease inhibitors were added (final concentration given): 2 µg/ml leupeptin, 10 µM phosphoramidon, 40  
20 µg/ml bacitracin, 5 µg/ml aprotinin, 0.1 mM Pefabloc.

Coated beads were kept on ice until use. For each well, 0.145 ml of beads were added to Optiplate assay plates (Packard 6005190), followed by 0.002-0.004 ml of competitor and 0.05 ml of [<sup>125</sup>I]-human MCH (Phe<sup>13</sup>, Tyr<sup>19</sup> substituted; 30 pM). Binding reactions were allowed to proceed at room temperature for 3 hours.  
25 Quantitation was performed by scintillation counting (TopCount device).

#### Example 4: MCH Activity

The activity of different MCH analogs was measured using the procedures described in Examples 2 and 3 above. Tables 1-7 illustrate the activity of  
30 different truncated MCH analogs and mammalian MCH (SEQ. ID. NO. 11). Figure 1 illustrates the results of replacing different amino acids of mammalian MCH with alanine. Based on the guidance provided herein, additional MCH analogs active at the MCH receptor can be obtained.

35

TABLE 1

SEQ. ID. NO.	Binding Assay IC <sub>50</sub> (nM)	EC <sub>50</sub> (nM)	% Activation at 10 $\mu$ M
11	0.3	36	100
7	0.12	18	123
8	0.16	36	123
9	1.6	300	74
10	0.3		99
12	6.4	492	3
13	1.5		65.9
14	0.5		62.2

IC<sub>50</sub> was determined using a SPA based assay.

EC<sub>50</sub> (nM) and % Activation at 10  $\mu$ M were determined using aequorin functional assays.

5

Table 2 illustrates the affect of different D-amino acids.

TABLE 2

Ac-Arg <sup>6</sup> -Cys <sup>7</sup> -Met <sup>8</sup> -Leu <sup>9</sup> -Gly <sup>10</sup> -Arg <sup>11</sup> -Val <sup>12</sup> -Tyr <sup>13</sup> -Arg <sup>14</sup> -Pro <sup>15</sup> -Cys <sup>16</sup> -NH <sub>2</sub>				
SEQ. ID. NO.	Compound	Activity		
		IC <sub>50</sub> (nM)	EC <sub>50</sub> (nM)	Activation %
11		0.3	30.9	100
7		0.5	20	99
15	D-Arg <sup>6</sup>	0.46	45	86
16	D-Cys <sup>7</sup>	7.78	909	34
17	D-Met <sup>8</sup>	>1000	Inactive	
18	D-Leu <sup>9</sup>	1520	Inactive	
19	D-Arg <sup>11</sup>	>1000	Inactive	
20	D-Val <sup>12</sup>	381	Inactive	
21	D-Tyr <sup>13</sup>	>1000	Inactive	
22	D-Arg <sup>14</sup>	368	Inactive	
23	D-Pro <sup>15</sup>	584	Inactive	
24	D-Cys <sup>16</sup>	0.8	133	76

10

Table 3 illustrates the effect of different N-methyl-amino acids.

TABLE 3

Ac-Arg <sup>6</sup> -Cys <sup>7</sup> -Met <sup>8</sup> -Leu <sup>9</sup> -Gly <sup>10</sup> -Arg <sup>11</sup> -Val <sup>12</sup> -Tyr <sup>13</sup> -Arg <sup>14</sup> -Pro <sup>15</sup> -Cys <sup>16</sup> -NH <sub>2</sub>				
SEQ. ID. NO.	Compound	Activity		
		IC <sub>50</sub> (nM)	EC <sub>50</sub> (nM)	Activation %
11		0.3	30.9	100
7		1.4	20	99
25	N-Me-Nle <sup>8</sup>	0.16	20	110
26	N-Me-Leu <sup>9</sup>	10%@1	>10000	3
27	Sar <sup>10</sup>	2.3	140	95
28	N-Me-Arg <sup>11</sup>	43	10	110
29	N-Me-Arg <sup>14</sup>	643	>1000	
30	Sar <sup>15</sup>	0.36	25	113

Table 4 illustrates the affect of different alterations to position 6 of the  
 5 SEQ. ID. NO. 7 MCH analog.

TABLE 4

X <sup>6</sup> -Cys <sup>7</sup> -Met <sup>8</sup> -Leu <sup>9</sup> -Gly <sup>10</sup> -Arg <sup>11</sup> -Val <sup>12</sup> -Tyr <sup>13</sup> -Arg <sup>14</sup> -Pro <sup>15</sup> -Cys <sup>16</sup> -NH <sub>2</sub>				
SEQ. ID. NO.	Position 6 modification	Activity		
		IC <sub>50</sub> (nM)	EC <sub>50</sub> (nM)	Activation %
11		0.3	30.9	100
7		1.4	20	99
31	Ac-Ala	27	114	135
32	Ac-Nle	40	117	107
33	Ac-Pro	3.4	59	133
34	Ac-Asn	2.6	150	96
35	Ac-Ser	4.5	207	120
36	Ac-Glu	19	935	113
37	H	12	809	120
38	Ac	1.6	144	82
39	Arg	0.13	14	106
40	Δ NH <sub>2</sub> -Arg	0.48	38.5	49
41	Ac-D-Arg	0.46	45	86
42	Ac-D-Nle	1.2	110	97
43	Ac-D-Pro	0.82	60	96
44	Ac-D-Asn	3	340	94
45	Ac-D-Ser	2.3	170	93
46	Ac-D-Glu	8	820	85

Table 5 illustrates the affect of different alterations to position 10 of the SEQ. ID. NO. 7 MCH analog.

TABLE 5

5

Ac-Arg <sup>6</sup> -Cys <sup>7</sup> -Met <sup>8</sup> -Leu <sup>9</sup> -X <sup>10</sup> -Arg <sup>11</sup> -Val <sup>12</sup> -Tyr <sup>13</sup> -Arg <sup>14</sup> -Pro <sup>15</sup> -Cys <sup>16</sup> -NH <sub>2</sub>				
SEQ. ID. NO.	Position 10 modification	Activity		
		IC <sub>50</sub> (nM)	EC <sub>50</sub> (nM)	Activation %
11		0.3	30.9	100
7		0.5	20	99
47	Ala	0.59	31	104
48	Leu	0.06	23	106
49	Nle	0.04	15	106
50	Pro	700	519	4
51	Asn	0.23	23	106
52	Ser	0.32	65	104
53	Lys	110	4500	25
54	Glu	190	> 10000	12
55	D-Leu	16	750	23
56	D-Nle	2.4	215	33
57	D-Pro	1.2	190	90
58	D-Glu	40% @ 1	>10000	
59	D-Lys	>1000	>10000	
60	β-Ala	390	>1000	3.2
61	γ-Abu	2.1	30.6	101

Table 6 illustrates the affect of different alterations to position 13 of the SEQ. ID. NO. 7 MCH analog.

TABLE 6

Ac-Arg <sup>6</sup> -Cys <sup>7</sup> -Met <sup>8</sup> -Leu <sup>9</sup> -Gly <sup>10</sup> -Arg <sup>11</sup> -Val <sup>12</sup> -Tyr <sup>13</sup> -Arg <sup>14</sup> -Pro <sup>15</sup> -Cys <sup>16</sup> -NH <sub>2</sub>				
SEQ. ID. NO.	Position 13 modification	Activity		
		IC <sub>50</sub> (nM)	EC <sub>50</sub> (nM)	Activation %
11		0.3	30.9	100
7		1.4	20	99
62	Phe	1	46	96
63	Trp	3.8	890	83
64	His	13.1	3400	66
65	(2)Nal	0.15	54	105
66	Phe(pF)	0.6	108	98
67	Phe(pNH <sub>2</sub> )	3.2	610	88
68	Phe(pCOOH)	>1000	>10000	
69	Cha	0.09	122	93

Table 7 illustrates the affect of some alteration combinations and some alterations to position 8 of the SEQ. ID. NO. 7 MCH analog.

TABLE 7


Ac-Arg <sup>6</sup> -Cys <sup>7</sup> -Met <sup>8</sup> -Leu <sup>9</sup> -Gly <sup>10</sup> -Arg <sup>11</sup> -Val <sup>12</sup> -Tyr <sup>13</sup> -Arg <sup>14</sup> -Pro <sup>15</sup> -Cys <sup>16</sup> -NH <sub>2</sub>				
SEQ. ID. NO.	Compound	Activity		
		IC <sub>50</sub> (nM)	EC <sub>50</sub> (nM)	Activation %
11		0.3	30.9	100
7		1.4	20	99
70	Ava <sup>9,10</sup>	3.7	587	82
71	D-Arg <sup>6</sup> ,Ava <sup>9,10</sup>	3.7	1080	72
72	Ava <sup>14,15</sup>	6.2	406	75
73	D-Arg <sup>6</sup> ,Ava <sup>14,15</sup>	19.5	1300	28
74	D-Pro <sup>10</sup> ,Ava <sup>14,15</sup>	700	1530	3
75	ΔArg <sup>6</sup> ,Ava <sup>14,15</sup>	250	>10000	3
76	Ava <sup>9,10</sup> ,Ava <sup>14,15</sup>	50	>10000	3
77	Nle <sup>8</sup>	0.5	44	105
78	ΔArg <sup>6</sup> ,D-Nle <sup>10</sup>	25	72	4

Other embodiments are within the following claims. While several embodiments have been shown and described, various modifications may be made without departing from the spirit and scope of the present invention.

## WHAT IS CLAIMED IS:

1. An optionally substituted peptide having the structure:

5



10 herein X<sup>1</sup> is an optionally present amino acid that, if present, is either alanine, valine, leucine, isoleucine, proline, tryptophan, phenylalanine, methionine, glycine, serine, threonine, tyrosine, cysteine, asparagine, glutamine, lysine, arginine, histidine, aspartic acid, or glutamic acid, or a derivative thereof;

15 X<sup>2</sup> is an optionally present amino acid that, if present, is either alanine, valine, leucine, isoleucine, proline, tryptophan, phenylalanine, methionine, glycine, serine, threonine, tyrosine, cysteine, asparagine, glutamine, lysine, arginine, histidine, aspartic acid, or glutamic acid, or a derivative thereof;

20 X<sup>3</sup> is an optionally present amino acid that, if present, is either alanine, valine, leucine, isoleucine, proline, tryptophan, phenylalanine, methionine, glycine, serine, threonine, tyrosine, cysteine, asparagine, glutamine, lysine, arginine, histidine, aspartic acid or glutamic acid, or a derivative thereof;

X<sup>4</sup> is an optionally present amino acid that, if present, is either alanine, valine, leucine, isoleucine, proline, tryptophan, phenylalanine, methionine, glycine, serine, threonine, tyrosine, cysteine, asparagine, glutamine, lysine, arginine, histidine, aspartic acid, glutamic acid, or norleucine, or a derivative thereof;

25 X<sup>5</sup> is an optionally present amino acid that, if present, is either alanine, valine, leucine, isoleucine, proline, tryptophan, phenylalanine, methionine, glycine, serine, threonine, tyrosine, cysteine, asparagine, glutamine, lysine, arginine, histidine, aspartic acid or glutamic acid, or a derivative thereof;

30 X<sup>6</sup> is an optionally present amino acid that, if present is either arginine, alanine, leucine, glycine, lysine, proline, asparagine, serine, histidine, nitroarginine, norleucine, or des-amino-arginine, or a derivative thereof,

X<sup>7</sup> is either cysteine, homocysteine, or penicillamine, or a derivative thereof;

35 X<sup>8</sup> is either methionine, norleucine, leucine, isoleucine, valine, methioninesulfoxide, or methioninesulfone, or a derivative thereof;

X<sup>9</sup> is either leucine, isoleucine, valine, alanine, methionine, or 5-aminopentanoic acid, or a derivative thereof;

X<sup>10</sup> is either glycine, alanine, leucine, norleucine, cyclohexylalanine, 5-aminopentanoic acid, asparagine, serine, sarcosine, isobutyric, or gamma-aminobutyric acid, or a derivative thereof;

X<sup>11</sup> is either arginine, lysine, citrulline, histidine, or nitroarginine, or a derivative thereof;

X<sup>12</sup> is either valine, leucine, isoleucine, alanine, or methionine, or a derivative thereof;

X<sup>13</sup> is either phenylalanine, tyrosine, D-(*p*-benzoylphenylalanine), tryptophan, (1')- and (2')-naphthylalanine, cyclohexylalanine, or mono and multi-substituted phenylalanine wherein each substituent is independently selected from the group consisting of O-alkyl, alkyl, OH, NO<sub>2</sub>, NH<sub>2</sub>, F, I, and Br; or a derivative thereof;

X<sup>14</sup> is either arginine, lysine, histidine, norarginine, or 5-aminopentanoic acid or a derivative thereof;

X<sup>15</sup> is either proline, alanine, valine, leucine, isoleucine, methionine, sarcosine, or 5-aminopentanoic acid, or a derivative thereof;

X<sup>16</sup> is either cysteine, homocysteine, or penicillamine, or a derivative thereof;

X<sup>17</sup> is an optionally present amino acid that, if present, is either alanine, valine, leucine, isoleucine, proline, tryptophan, phenylalanine, methionine, glycine, serine, threonine, tyrosine, cysteine, asparagine, glutamine, lysine, arginine, histidine, aspartic acid or glutamic acid, or a derivative thereof;

Z<sup>1</sup> is an optionally present protecting group that, if present, is covalently joined to the N-terminal amino group;

Z<sup>2</sup> is an optionally present protecting group that, if present, is covalently joined to the C-terminal carboxy group;

or a labeled derivative of said peptide;

or a pharmaceutically acceptable salt of said peptide or of said labeled derivative.

2. The peptide of claim 1, wherein X<sup>1</sup>, X<sup>2</sup>, X<sup>3</sup>, X<sup>4</sup>, and X<sup>5</sup> are not present; X<sup>6</sup> is arginine; and X<sup>17</sup> is tyrosine or tryptophan.

3. The peptide of claim 1, wherein X<sup>1</sup>, X<sup>2</sup>, X<sup>3</sup>, X<sup>4</sup>, X<sup>5</sup>, and X<sup>17</sup> are not present; and X<sup>6</sup> is arginine.

5 4. The peptide of claim 3, wherein Z<sup>1</sup> is -C(O)CH<sub>3</sub> and Z<sup>2</sup> is -NH<sub>2</sub>.

5. The peptide of claim 1, wherein said peptide is either SEQ. ID. NO. 7, SEQ. ID. NO. 8, SEQ. ID. NO. 9, SEQ. ID. NO. 10, or  
10 a pharmaceutically acceptable salt thereof.

6. The peptide of claim 1, wherein said peptide is either SEQ. ID. NO. 7, SEQ. ID. NO. 8, or a pharmaceutically acceptable salt thereof.

15 7. The peptide of claim 1, wherein X<sup>1</sup>, X<sup>2</sup>, X<sup>3</sup>, X<sup>4</sup>, X<sup>5</sup> and X<sup>17</sup> are not present;  
X<sup>6</sup> is either arginine, D-arginine, D-norleucine, D-proline, D-serine, or D-asparagine;  
X<sup>7</sup> is cysteine;  
X<sup>8</sup> is either methionine, norleucine, or N-methyl norleucine;  
20 X<sup>9</sup> is leucine;  
X<sup>10</sup> is either glycine, alanine, leucine, norleucine, asparagine, serine, D-norleucine, D-proline, gamma-aminobutyric acid, or sarcosine;  
X<sup>11</sup> is arginine;  
X<sup>12</sup> is valine;  
25 X<sup>13</sup> is phenylalanine, (2')naphthylalanine, p-fluoro-phenylalanine, tyrosine, or cyclohexylalanine;  
X<sup>14</sup> is arginine;  
X<sup>15</sup> is either proline or sarcosine; and  
X<sup>16</sup> is either cysteine or D-cysteine.

30

8. The peptide of claim 1, wherein said peptide consists of a sequence selected from the group consisting of: 7, 8, 10, 15, 24, 25, 27, 28, 30-49, 51, 52, 56, 57, 61, 62, 63, 65-67, 69-72, and 77.

9. A method of screening for a compound able to bind a MCH receptor comprising the step of measuring the ability of said compound to effect binding of the peptide of any one of claims 1-8 to either said receptor, a fragment of said receptor comprising a MCH binding site, a polypeptide comprising said fragment, or a derivative of said polypeptide.

10. The method of claim 9, wherein said method measures the ability of said peptide to bind to said receptor or said fragment thereof.

11. The method of claim 10, wherein said peptide is radiolabeled.

12. The method of claim 9, wherein said peptide is a radiolabeled derivative of SEQ. ID. NO. 8 or a pharmaceutically acceptable salt thereof.

13. A method for increasing weight in a subject comprising the step of administering to said subject an effective amount of the peptide of any one of claims 1-8 to produce a weight increase.

14. A method for increasing appetite in a subject comprising the step of administering to said subject an effective amount of the peptide of any one of claims 1-8 to produce an appetite increase.

15. A method for measuring the ability of a compound to decrease weight or appetite in a subject comprising the steps of:

- a) administering to said subject an effective amount of the peptide of any one of claims 1-8 to produce a weight increase or appetite increase,
- b) administering said compound to said subject, and
- c) measuring the change in weight or appetite of said subject.

16. The method of claim 15, wherein said subject is either a rat or a mouse.

1/1

ALANINE SCAN OF HUMAN MELANIN-CONCENTRATING HORMONE (hMCH)

	1	3	5	*	9	11	13	15	*	17	19								
Asp Phe Asp Met Leu Arg Cys Met Leu Gly Arg Val Tyr Arg Pro Cys Trp Gln Val ACID																			
BINDING IC <sub>50</sub> (nM)	0.8	0.2	0.49	0.49	0.33	4.5	-	71	0.9	-	169	0.9	253	1.4	1.9	-	0.2	1.1	0.3
FUNCTION EC <sub>50</sub> (nM)	ND	ND	ND	ND	ND	ND	-	ND	ND	-	ND	ND	ND	ND	ND	-	ND	ND	ND
% ACTIVATION (100 nM)	97.7	96.3	97.7	95.7	90.8	11.4	-	0	86.5	-	3	82.9	0	90.4	76.3	-	100	78.3	94.4

hMCH

BINDING ASSAY<sup>+</sup>, IC<sub>50</sub> = 0.3nM

AEQUORIN FUNCTIONAL ASSAY<sup>++</sup>, EC<sub>50</sub> = 36nM, 100% ACTIVATION AT 10μM

+ SPA ASSAY: INHIBITION OF (<sup>125</sup>I-F13, Y19-hMCH BINDING TO THE CLONED HUMAN MCH RECEPTOR (COS-7 CELLS)  
 ++ FOR AEQUORIN FUNCTIONAL ASSAY, 100% ACTIVATION IS THE BIOLUMINESCENCE VALUE OBTAINED WITH 100 nM MCH.  
 \* SITES OF CYCLIZATION (S-S)  
 ND-NOT DONE

FIG.1

## SEQUENCE LISTING

&lt;110&gt; Merck &amp; Co., Inc.

&lt;120&gt; MELANIN-CONCENTRATING HORMONE ANALOGS

&lt;130&gt; 20590Y PCT

&lt;150&gt; 60/179,967

&lt;151&gt; 2000-02-03

&lt;160&gt; 78

&lt;170&gt; FastSEQ for Windows Version 4.0

&lt;210&gt; 1

&lt;211&gt; 1062

&lt;212&gt; DNA

&lt;213&gt; Human

```

<400> 1
atggacctgg aagcctcgct gctgcccact ggtcccaacg ccagcaacac ctctgatggc      60
cccgataacc tcacttcggc aggatcacct cctcgcacgg ggagcatctc ctacatcaac      120
atcatcatgc cttcggtgtt cggcaccatc tgcctcctgg gcatcatcgg gaactccacg      180
gtcatcttcg cggtcgtgaa gaagtccaag ctgcaactgg gcaacaacgt ccccgcacac      240
ttcatcatca acctctcggt agtagatctc ctctttctcc tgggcatgcc ctcatgatc      300
caccagctca tgggcaatgg ggtgtggcac tttggggaga ccatgtgcac cctcatcacg      360
gccatggatg ccaatagtca gtaccaccagc acctacatcc tgaccgccat ggccattgac      420
cgctacctgg ccaactgtcca ccccatctct tccacgaagt tccggaagcc ctctgtggcc      480
accctgggtg tctgctctct gtgggccctc tccttcatca gcatcaccac tgtgtggctg      540
tatgccagac tcacccctt cccaggaggt gcagtgggct gcggcatacg cctgcccac      600
ccagacactg acctctactg gtaccacctg taccagtttt tcctggcctt tgccctgcct      660
tttgtggtca tcacagccgc atactgtagg atcctgcagc gcatgacgtc ctcagtggcc      720
ccgcctcccc agcgcagcat cggctgcgg acaaagaggg tgaccgcgac agccatcgcc      780
atctgtctgg tcttctttgt gtgtgtggca cctactatg tgctacagct gaccagttg      840
tccatcagcc gcccgacct cactttgttc tacttataca atgcgccat cagcttgggc      900
tatgccaaca gctgcctcaa cccctttgtg tacatcgtgc tctgtgagac gttccgcaa      960
cgcttggttc tgtcggtgaa gcctgcagcc caggggcagc ttcgcgctgt cagcaacgct     1020
cagacggctg acgaggagag gacagaaagc aaaggcacct ga                               1062

```

&lt;210&gt; 2

&lt;211&gt; 1254

&lt;212&gt; DNA

&lt;213&gt; Human

```

<400> 2
atgaagaagg gagtggggag ggcagttggg cttggaggcg gcagcggctg ccaggctacg      60
gaggaagacc cccttcccaa ctgcggggct tgcgctccgg gacaagggtg caggcgctgg      120
aggctgccgc agcctgcgtg ggtggagggg agctcagctc ggttgtggga gcaggcgacc      180
ggcactggct ggatggacct ggaagcctcg ctgctgcca ctggtcccaa cgccagcaac      240
acctctgatg gccccgataa cctcacttcg gcaggatcac ctctcgcac ggggagcatc      300
tcctacatca acatcatcat gccttcgggtg ttcggcacca tctgcctcct gggcatcatc      360
gggaactcca cggatcatctt cgcggtcgtg aagaagtcca agctgcactg gtgcaacaac      420
gtccccgaca tcttcatcat caacctctcg gtagtagatc tcctctttct cctgggcatg      480
cccttcatga tccaccagct catgggcaat ggggtgtggc actttgggga gaccatgtgc      540
accctcatca cggccatgga tgccaatagt cagttcacca gcacctacat cctgaccgcc      600
atggccattg accgctacct ggccactgtc caccctatct cttccacgaa gttccggaag      660
ccctctgttg ccacctggt gatctgcctc ctgtgggccc tctccttcat cagcatcacc      720
cctgtgtggc tgtatgccag actcatcccc ttcccaggag gtgcagtggg ctgcggcata      780
cgcttgccca acccagacac tgacctctac tgggtcaccc tgtaccagtt tttcctggcc      840

```

tttgccctgc	ctttttgtggt	catcacagcc	gcatacgtga	ggatcctgca	gcgcatgacg	900
tcctcagtgg	ccccgcctc	ccagcgcagc	atccgggtgc	ggacaaagag	ggtgacccgc	960
acagccatcg	ccatctgtct	ggtcttcttt	gtgtgctggg	caccctacta	tgtgctacag	1020
ctgaccacgt	tgtccatcag	cgcgccgacc	ctcacctttg	tctacttata	caatgcggcc	1080
atcagcttgg	gctatgccaa	cagctgcctc	aacccctttg	tgtacatcgt	gctctgtgag	1140
acgttcgcga	aacgcttgg	cctgtcgggt	aagcctgcag	cccaggggca	gcttcgcgct	1200
gtcagcaacg	ctcagacggc	tgacgaggag	aggacagaaa	gcaaaggcac	ctga	1254

<210> 3  
 <211> 1269  
 <212> DNA  
 <213> Human

<400> 3							
atgtcagtgg	gagccatgaa	gaagggagtg	gggagggcag	ttgggcttgg	aggcggcagc	60	
ggctgccagg	ctacggagga	agacccccctt	cccaactgcg	gggcttgccg	tccgggacaa	120	
ggtggcaggc	gctggaggct	gcccgcagcct	gcgtgggtgg	aggggagctc	agctcgggtg	180	
tgggagcagg	cgaccggcac	tggttgatg	gacctggaag	cctcgctgct	gccactggt	240	
cccaacgcca	gcaacacctc	tgatggcccc	gataacctca	cttcggcagg	atcacctcct	300	
cgcacgggga	gcatctccta	catcaacatc	atcatgcctt	cggtgttcgg	caccatctgc	360	
ctcctgggca	tcacgggaa	ctccacggtc	atcttcgcgg	tcgtgaagaa	gtccaagctg	420	
cactgggtgc	acaacgtccc	cgacatcttc	atcatcaacc	tctcggtagt	agatctcctc	480	
ttctccttgg	gcatgccctt	catgatccac	cagctcatgg	gcaatggggg	gtggcacttt	540	
ggggagacca	tgtgcacct	catcacggcc	atggatgcca	atagtcagtt	caccagcacc	600	
tacatcctga	ccgcatggc	cattgaccgc	tacctggcca	ctgtccaccc	catctcttcc	660	
acgaagtcc	ggaagccctc	tgtggccacc	ctgggtgatc	gcctcctgtg	ggccctctcc	720	
ttcatcagca	tcaccctgt	gtggctgtat	gccagactca	tccccctccc	aggaggtgca	780	
gtgggctgcg	gcatacgct	gcccacccca	gacactgacc	tctactgggt	cacctgttac	840	
cagtttttcc	tggcctttgc	cctgcctttt	gtggctcatca	cagccgcata	cgtaggagtc	900	
ctgcagcgca	tgacgtcctc	agtggccccc	gcctcccagc	gcagcatccg	gcggcgga	960	
aagaggggtga	cccgacagc	catcgccatc	tgtctggtct	tctttgtgtg	ctgggcaccc	1020	
tactatgtgc	tacagctgac	ccagttgtcc	atcagccgcc	cgacctcac	ctttgtctac	1080	
ttatacaatg	cggccatcag	cttgggctat	gccaacagct	gcctcaaccc	ctttgtgtac	1140	
atcgtgctct	gtgagacgtt	ccgcaaacgc	ttggctcctgt	cggtgaagcc	tgcagcccag	1200	
gggcagcttc	gcgtgtcag	caacgctcag	acggctgacg	aggagaggac	agaaagcaaa	1260	
ggcacctga						1269	

<210> 4  
 <211> 353  
 <212> PRT  
 <213> Human

<400> 4														
Met Asp Leu Glu Ala Ser Leu Leu Pro Thr Gly Pro Asn Ala Ser Asn														
1				5				10					15	
Thr Ser Asp Gly Pro Asp Asn Leu Thr Ser Ala Gly Ser Pro Pro Arg														
			20				25					30		
Thr Gly Ser Ile Ser Tyr Ile Asn Ile Ile Met Pro Ser Val Phe Gly														
			35				40					45		
Thr Ile Cys Leu Leu Gly Ile Ile Gly Asn Ser Thr Val Ile Phe Ala														
			50				55					60		
Val Val Lys Lys Ser Lys Leu His Trp Cys Asn Asn Val Pro Asp Ile														
65						70			75				80	
Phe Ile Ile Asn Leu Ser Val Val Asp Leu Leu Phe Leu Leu Gly Met														
			85				90						95	
Pro Phe Met Ile His Gln Leu Met Gly Asn Gly Val Trp His Phe Gly														
			100				105						110	
Glu Thr Met Cys Thr Leu Ile Thr Ala Met Asp Ala Asn Ser Gln Phe														
			115				120						125	
Thr Ser Thr Tyr Ile Leu Thr Ala Met Ala Ile Asp Arg Tyr Leu Ala														
			130				135						140	

Thr Val His Pro Ile Ser Ser Thr Lys Phe Arg Lys Pro Ser Val Ala  
 145 150 155 160  
 Thr Leu Val Ile Cys Leu Leu Trp Ala Leu Ser Phe Ile Ser Ile Thr  
 165 170 175  
 Pro Val Trp Leu Tyr Ala Arg Leu Ile Pro Phe Pro Gly Gly Ala Val  
 180 185 190  
 Gly Cys Gly Ile Arg Leu Pro Asn Pro Asp Thr Asp Leu Tyr Trp Phe  
 195 200 205  
 Thr Leu Tyr Gln Phe Phe Leu Ala Phe Ala Leu Pro Phe Val Val Ile  
 210 215 220  
 Thr Ala Ala Tyr Val Arg Ile Leu Gln Arg Met Thr Ser Ser Val Ala  
 225 230 235 240  
 Pro Ala Ser Gln Arg Ser Ile Arg Leu Arg Thr Lys Arg Val Thr Arg  
 245 250 255  
 Thr Ala Ile Ala Ile Cys Leu Val Phe Phe Val Cys Trp Ala Pro Tyr  
 260 265 270  
 Tyr Val Leu Gln Leu Thr Gln Leu Ser Ile Ser Arg Pro Thr Leu Thr  
 275 280 285  
 Phe Val Tyr Leu Tyr Asn Ala Ala Ile Ser Leu Gly Tyr Ala Asn Ser  
 290 295 300  
 Cys Leu Asn Pro Phe Val Tyr Ile Val Leu Cys Glu Thr Phe Arg Lys  
 305 310 315 320  
 Arg Leu Val Leu Ser Val Lys Pro Ala Ala Gln Gly Gln Leu Arg Ala  
 325 330 335  
 Val Ser Asn Ala Gln Thr Ala Asp Glu Glu Arg Thr Glu Ser Lys Gly  
 340 345 350  
 Thr

<210> 5  
 <211> 417  
 <212> PRT  
 <213> Human

<400> 5  
 Met Lys Lys Gly Val Gly Arg Ala Val Gly Leu Gly Gly Gly Ser Gly  
 1 5 10 15  
 Cys Gln Ala Thr Glu Glu Asp Pro Leu Pro Asn Cys Gly Ala Cys Ala  
 20 25 30  
 Pro Gly Gln Gly Gly Arg Arg Trp Arg Leu Pro Gln Pro Ala Trp Val  
 35 40 45  
 Glu Gly Ser Ser Ala Arg Leu Trp Glu Gln Ala Thr Gly Thr Gly Trp  
 50 55 60  
 Met Asp Leu Glu Ala Ser Leu Leu Pro Thr Gly Pro Asn Ala Ser Asn  
 65 70 75 80  
 Thr Ser Asp Gly Pro Asp Asn Leu Thr Ser Ala Gly Ser Pro Pro Arg  
 85 90 95  
 Thr Gly Ser Ile Ser Tyr Ile Asn Ile Ile Met Pro Ser Val Phe Gly  
 100 105 110  
 Thr Ile Cys Leu Leu Gly Ile Ile Gly Asn Ser Thr Val Ile Phe Ala  
 115 120 125  
 Val Val Lys Lys Ser Lys Leu His Trp Cys Asn Asn Val Pro Asp Ile  
 130 135 140  
 Phe Ile Ile Asn Leu Ser Val Val Asp Leu Leu Phe Leu Leu Gly Met  
 145 150 155 160  
 Pro Phe Met Ile His Gln Leu Met Gly Asn Gly Val Trp His Phe Gly  
 165 170 175  
 Glu Thr Met Cys Thr Leu Ile Thr Ala Met Asp Ala Asn Ser Gln Phe  
 180 185 190  
 Thr Ser Thr Tyr Ile Leu Thr Ala Met Ala Ile Asp Arg Tyr Leu Ala  
 195 200 205

Thr Val His Pro Ile Ser Ser Thr Lys Phe Arg Lys Pro Ser Val Ala  
 210 215 220  
 Thr Leu Val Ile Cys Leu Leu Trp Ala Leu Ser Phe Ile Ser Ile Thr  
 225 230 235 240  
 Pro Val Trp Leu Tyr Ala Arg Leu Ile Pro Phe Pro Gly Gly Ala Val  
 245 250 255  
 Gly Cys Gly Ile Arg Leu Pro Asn Pro Asp Thr Asp Leu Tyr Trp Phe  
 260 265 270  
 Thr Leu Tyr Gln Phe Phe Leu Ala Phe Ala Leu Pro Phe Val Val Ile  
 275 280 285  
 Thr Ala Ala Tyr Val Arg Ile Leu Gln Arg Met Thr Ser Ser Val Ala  
 290 295 300  
 Pro Ala Ser Gln Arg Ser Ile Arg Leu Arg Thr Lys Arg Val Thr Arg  
 305 310 315 320  
 Thr Ala Ile Ala Ile Cys Leu Val Phe Phe Val Cys Trp Ala Pro Tyr  
 325 330 335  
 Tyr Val Leu Gln Leu Thr Gln Leu Ser Ile Ser Arg Pro Thr Leu Thr  
 340 345 350  
 Phe Val Tyr Leu Tyr Asn Ala Ala Ile Ser Leu Gly Tyr Ala Asn Ser  
 355 360 365  
 Cys Leu Asn Pro Phe Val Tyr Ile Val Leu Cys Glu Thr Phe Arg Lys  
 370 375 380  
 Arg Leu Val Leu Ser Val Lys Pro Ala Ala Gln Gly Gln Leu Arg Ala  
 385 390 395 400  
 Val Ser Asn Ala Gln Thr Ala Asp Glu Glu Arg Thr Glu Ser Lys Gly  
 405 410 415  
 Thr

<210> 6  
 <211> 422  
 <212> PRT  
 <213> Human

<400> 6  
 Met Ser Val Gly Ala Met Lys Lys Gly Val Gly Arg Ala Val Gly Leu  
 1 5 10 15  
 Gly Gly Gly Ser Gly Cys Gln Ala Thr Glu Glu Asp Pro Leu Pro Asn  
 20 25 30  
 Cys Gly Ala Cys Ala Pro Gly Gln Gly Gly Arg Arg Trp Arg Leu Pro  
 35 40 45  
 Gln Pro Ala Trp Val Glu Gly Ser Ser Ala Arg Leu Trp Glu Gln Ala  
 50 55 60  
 Thr Gly Thr Gly Trp Met Asp Leu Glu Ala Ser Leu Leu Pro Thr Gly  
 65 70 75 80  
 Pro Asn Ala Ser Asn Thr Ser Asp Gly Pro Asp Asn Leu Thr Ser Ala  
 85 90 95  
 Gly Ser Pro Pro Arg Thr Gly Ser Ile Ser Tyr Ile Asn Ile Ile Met  
 100 105 110  
 Pro Ser Val Phe Gly Thr Ile Cys Leu Leu Gly Ile Ile Gly Asn Ser  
 115 120 125  
 Thr Val Ile Phe Ala Val Val Lys Lys Ser Lys Leu His Trp Cys Asn  
 130 135 140  
 Asn Val Pro Asp Ile Phe Ile Ile Asn Leu Ser Val Val Asp Leu Leu  
 145 150 155 160  
 Phe Leu Leu Gly Met Pro Phe Met Ile His Gln Leu Met Gly Asn Gly  
 165 170 175  
 Val Trp His Phe Gly Glu Thr Met Cys Thr Leu Ile Thr Ala Met Asp  
 180 185 190  
 Ala Asn Ser Gln Phe Thr Ser Thr Tyr Ile Leu Thr Ala Met Ala Ile  
 195 200 205

Asp Arg Tyr Leu Ala Thr Val His Pro Ile Ser Ser Thr Lys Phe Arg  
 210 215 220  
 Lys Pro Ser Val Ala Thr Leu Val Ile Cys Leu Leu Trp Ala Leu Ser  
 225 230 235 240  
 Phe Ile Ser Ile Thr Pro Val Trp Leu Tyr Ala Arg Leu Ile Pro Phe  
 245 250 255  
 Pro Gly Gly Ala Val Gly Cys Gly Ile Arg Leu Pro Asn Pro Asp Thr  
 260 265 270  
 Asp Leu Tyr Trp Phe Thr Leu Tyr Gln Phe Phe Leu Ala Phe Ala Leu  
 275 280 285  
 Pro Phe Val Val Ile Thr Ala Tyr Val Arg Ile Leu Gln Arg Met  
 290 295 300  
 Thr Ser Ser Val Ala Pro Ala Ser Gln Arg Ser Ile Arg Leu Arg Thr  
 305 310 315 320  
 Lys Arg Val Thr Arg Thr Ala Ile Ala Ile Cys Leu Val Phe Phe Val  
 325 330 335  
 Cys Trp Ala Pro Tyr Tyr Val Leu Gln Leu Thr Gln Leu Ser Ile Ser  
 340 345 350  
 Arg Pro Thr Leu Thr Phe Val Tyr Leu Tyr Asn Ala Ala Ile Ser Leu  
 355 360 365  
 Gly Tyr Ala Asn Ser Cys Leu Asn Pro Phe Val Tyr Ile Val Leu Cys  
 370 375 380  
 Glu Thr Phe Arg Lys Arg Leu Val Leu Ser Val Lys Pro Ala Ala Gln  
 385 390 395 400  
 Gly Gln Leu Arg Ala Val Ser Asn Ala Gln Thr Ala Asp Glu Glu Arg  
 405 410 415  
 Thr Glu Ser Lys Gly Thr  
 420

<210> 7  
 <211> 11  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <221> ACETYLATION  
 <222> (1)...(1)

<221> AMIDATION  
 <222> (11)...(11)

<221> DISULFID  
 <222> (2)...(11)

<223> MCH Analog

<400> 7  
 Arg Cys Met Leu Gly Arg Val Tyr Arg Pro Cys  
 1 5 10

<210> 8  
 <211> 12  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <221> ACETYLATION  
 <222> (1)...(1)

<221> AMIDATION  
 <222> (12)...(12)

<221> DISULFID  
<222> (2)...(11)  
<223> MCH Analog  
<400> 8  
Arg Cys Met Leu Gly Arg Val Phe Arg Pro Cys Tyr  
1 5 10  
<210> 9  
<211> 10  
<212> PRT  
<213> Artificial Sequence  
<220>  
<221> ACETYLATION  
<222> (1)...(1)  
<221> AMIDATION  
<222> (10)...(10)  
<221> DISULFID  
<222> (1)...(10)  
<223> MCH Analog  
<400> 9  
Cys Met Leu Gly Arg Val Tyr Arg Pro Cys  
1 5 10  
<210> 10  
<211> 16  
<212> PRT  
<213> Artificial Sequence  
<220>  
<221> AMIDATION  
<222> (16)...(16)  
<221> DISULFID  
<222> (7)...(16)  
<223> MCH Analog  
<400> 10  
Asp Phe Asp Met Leu Arg Cys Met Leu Gly Arg Val Tyr Arg Pro Cys  
1 5 10 15  
<210> 11  
<211> 19  
<212> PRT  
<213> Human  
<220>  
<221> DISULFID  
<222> (7)...(16)  
<400> 11  
Asp Phe Asp Met Leu Arg Cys Met Leu Gly Arg Val Tyr Arg Pro Cys  
1 5 10 15  
Trp Gln Val

<210> 12  
<211> 13  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> MCH Analog

<221> DISULFID  
<222> (1)...(10)

<221> ACETYLTATION  
<222> (1)...(1)

<400> 12  
Cys Met Leu Gly Arg Val Tyr Arg Pro Cys Trp Gln Val  
1 5 10

<210> 13  
<211> 19  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> MCH Analog

<221> DISULFID  
<222> (7)...(16)

<221> MOD\_RES  
<222> (4)...(4)  
<223> Xaa = Norleucine

<221> MOD\_RES  
<222> (8)...(8)  
<223> Xaa = Norleucine

<400> 13  
Asp Phe Asp Xaa Leu Arg Cys Xaa Leu Gly Arg Val Tyr Arg Pro Cys  
1 5 10 15  
Trp Gln Val

<210> 14  
<211> 19  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> MCH Analog

<221> DISULFID  
<222> (7)...(16)

<400> 14  
Asp Phe Ala Met Leu Arg Cys Met Leu Gly Arg Val Phe Arg Pro Cys  
1 5 10 15  
Trp Gln Tyr

<210> 15

<211> 11  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> MCH Analogs

<221> DISULFID  
<222> (2)...(11)

<221> ACETYLATION  
<222> (1)...(1)

<221> MOD\_RES  
<222> (1)...(1)  
<223> Xaa = D-Arginine

<221> AMIDATION  
<222> (11)...(11)

<400> 15  
Xaa Cys Met Leu Gly Arg Val Tyr Arg Pro Cys  
1 5 10

<210> 16  
<211> 11  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> MCH Analog

<221> DISULFID  
<222> (2)...(11)

<221> MOD\_RES  
<222> (2)...(2)  
<223> Xaa = D-Cysteine

<221> ACETYLATION  
<222> (1)...(1)

<221> AMIDATION  
<222> (11)...(11)

<400> 16  
Arg Xaa Met Leu Gly Arg Val Tyr Arg Pro Cys  
1 5 10

<210> 17  
<211> 11  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> MCH Analog

<221> DISULFID  
<222> (2)...(11)

<221> MOD\_RES  
<222> (3)...(3)

<223> Xaa = D-Methionine

<221> ACETYLATION

<222> (1)...(1)

<221> AMIDATION

<222> (11)...(11)

<400> 17

Arg Cys Xaa Leu Gly Arg Val Tyr Arg Pro Cys  
1 5 10

<210> 18

<211> 11

<212> PRT

<213> Artificial Sequence

<220>

<223> MCH Analog

<221> DISULFID

<222> (2)...(11)

<221> MOD\_RES

<222> (4)...(4)

<223> Xaa = D-Leucine

<221> ACETYLATION

<222> (1)...(1)

<221> AMIDATION

<222> (11)...(11)

<400> 18

Arg Cys Met Xaa Gly Arg Val Tyr Arg Pro Cys  
1 5 10

<210> 19

<211> 11

<212> PRT

<213> Artificial Sequence

<220>

<223> MCH Analog

<221> DISULFID

<222> (2)...(11)

<221> MOD\_RES

<222> (6)...(6)

<223> Xaa = D-Arginine

<221> ACETYLATION

<222> (1)...(1)

<221> AMIDATION

<222> (11)...(11)

<400> 19

Arg Cys Met Leu Gly Xaa Val Tyr Arg Pro Cys  
1 5 10

<210> 20  
<211> 11  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> MCH Analog

<221> DISULFID  
<222> (2)...(11)

<221> MOD\_RES  
<222> (7)...(7)  
<223> Xaa = D-Valine

<221> ACETYLATION  
<222> (1)...(1)

<221> AMIDATION  
<222> (11)...(11)

<400> 20  
Arg Cys Met Leu Gly Arg Xaa Tyr Arg Pro Cys  
1 5 10

<210> 21  
<211> 11  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> MCH Analog

<221> DISULFID  
<222> (2)...(11)

<221> MOD\_RES  
<222> (8)...(8)  
<223> Xaa = D-Tyrosine

<221> ACETYLATION  
<222> (1)...(1)

<221> AMIDATION  
<222> (11)...(11)

<400> 21  
Arg Cys Met Leu Gly Arg Val Xaa Arg Pro Cys  
1 5 10

<210> 22  
<211> 11  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> MCH Analog

<221> ACETYLATION  
<222> (1)...(1)

<221> MOD\_RES

<222> (9)...(9)  
<223> Xaa = D-Arginine

<221> DISULFID  
<222> (2)...(11)

<221> AMIDATION  
<222> (11)...(11)

<400> 22  
Arg Cys Met Leu Gly Arg Val Tyr Xaa Pro Cys  
1 5 10

<210> 23  
<211> 11  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> MCH Analog

<221> ACETYLATION  
<222> (1)...(1)

<221> MOD\_RES  
<222> (10)...(10)  
<223> Xaa = D-Proline

<221> DISULFID  
<222> (2)...(11)

<221> AMIDATION  
<222> (11)...(11)

<400> 23  
Arg Cys Met Leu Gly Arg Val Tyr Arg Xaa Cys  
1 5 10

<210> 24  
<211> 11  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> MCH Analog

<221> ACETYLATION  
<222> (1)...(1)

<221> MOD\_RES  
<222> (11)...(11)  
<223> Xaa = D-Cysteine

<221> DISULFID  
<222> (2)...(11)

<221> AMIDATION  
<222> (11)...(11)

<400> 24  
Arg Cys Met Leu Gly Arg Val Tyr Arg Pro Xaa  
1 5 10

<210> 25  
<211> 11  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> MCH Analog

<221> ACETYLATION  
<222> (1)...(1)

<221> MOD\_RES  
<222> (3)...(3)  
<223> Xaa = N-Me-Norleucine

<221> DISULFID  
<222> (2)...(11)

<221> AMIDATION  
<222> (11)...(11)

<400> 25  
Arg Cys Xaa Leu Gly Arg Val Tyr Arg Pro Cys  
1 5 10

<210> 26  
<211> 11  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> MCH Analog

<221> ACETYLATION  
<222> (1)...(1)

<221> MOD\_RES  
<222> (4)...(4)  
<223> Xaa = N-Me-Leucine

<221> DISULFID  
<222> (2)...(11)

<221> AMIDATION  
<222> (11)...(11)

<400> 26  
Arg Cys Met Xaa Gly Arg Val Tyr Arg Pro Cys  
1 5 10

<210> 27  
<211> 11  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> MCH Analog

<221> ACETYLATION  
<222> (1)...(1)

<221> MOD\_RES  
<222> (5)...(5)  
<223> Xaa = MeGly

<221> DISULFID  
<222> (2)...(11)

<221> AMIDATION  
<222> (11)...(11)

<400> 27  
Arg Cys Met Leu Xaa Arg Val Tyr Arg Pro Cys  
1 5 10

<210> 28  
<211> 11  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> MCH Analog

<221> ACETYLTATION  
<222> (1)...(1)

<221> MOD\_RES  
<222> (6)...(6)  
<223> Xaa = N-Me-Arginine

<221> DISULFID  
<222> (2)...(11)

<221> AMIDATION  
<222> (11)...(11)

<400> 28  
Arg Cys Met Leu Gly Xaa Val Tyr Arg Pro Cys  
1 5 10

<210> 29  
<211> 11  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> MCH Analog

<221> ACETYLTATION  
<222> (1)...(1)

<221> MOD\_RES  
<222> (9)...(9)  
<223> Xaa = N-Me-Arginine

<221> DISULFID  
<222> (2)...(11)

<221> AMIDATION  
<222> (11)...(11)

<400> 29

Arg Cys Met Leu Gly Arg Val Tyr Xaa Pro Cys  
1 5 10

<210> 30  
<211> 11  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> MCH Analog  
  
<221> ACETYLATION  
<222> (1)...(1)

<221> MOD\_RES  
<222> (10)...(10)  
<223> Xaa = MeGly

<221> DISULFID  
<222> (2)...(11)

<221> AMIDATION  
<222> (11)...(11)

<400> 30  
Arg Cys Met Leu Gly Arg Val Tyr Arg Xaa Cys  
1 5 10

<210> 31  
<211> 11  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> MCH Analog  
  
<221> ACETYLATION  
<222> (1)...(1)

<221> DISULFID  
<222> (2)...(11)

<221> AMIDATION  
<222> (11)...(11)

<400> 31  
Ala Cys Met Leu Gly Arg Val Tyr Arg Pro Cys  
1 5 10

<210> 32  
<211> 11  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> MCH Analog

<221> MOD\_RES  
<222> (1)...(1)  
<223> Xaa = Norleucine

<221> DISULFID

<222> (2)...(11)

<221> AMIDATION  
<222> (11)...(11)

<221> ACETYLATION  
<222> (1)...(1)

<400> 32  
Xaa Cys Met Leu Gly Arg Val Tyr Arg Pro Cys  
1 5 10

<210> 33  
<211> 11  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> MCH Analog

<221> ACETYLATION  
<222> (1)...(1)

<221> DISULFID  
<222> (2)...(11)

<221> AMIDATION  
<222> (11)...(11)

<400> 33  
Pro Cys Met Leu Gly Arg Val Tyr Arg Pro Cys  
1 5 10

<210> 34  
<211> 11  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> MCH Analog

<221> ACETYLATION  
<222> (1)...(1)

<221> DISULFID  
<222> (2)...(11)

<221> AMIDATION  
<222> (11)...(11)

<400> 34  
Asn Cys Met Leu Gly Arg Val Tyr Arg Pro Cys  
1 5 10

<210> 35  
<211> 11  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> MCH Analog

<221> ACETYLATION  
<222> (1)...(1)

<221> DISULFID  
<222> (2)...(11)

<221> AMIDATION  
<222> (11)...(11)

<400> 35  
Ser Cys Met Leu Gly Arg Val Tyr Arg Pro Cys  
1 5 10

<210> 36  
<211> 11  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> MCH Analog

<221> ACETYLATION  
<222> (1)...(1)

<221> DISULFID  
<222> (2)...(11)

<221> AMIDATION  
<222> (11)...(11)

<400> 36  
Glu Cys Met Leu Gly Arg Val Tyr Arg Pro Cys  
1 5 10

<210> 37  
<211> 10  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> MCH Analog

<221> AMIDATION  
<222> (10)...(10)

<221> DISULFID  
<222> (1)...(10)

<400> 37  
Cys Met Leu Gly Arg Val Tyr Arg Pro Cys  
1 5 10

<210> 38  
<211> 10  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> MCH Analog

<221> ACETYLATION  
<222> (1)...(1)

<221> DISULFID  
<222> (1)...(10)

<221> AMIDATION  
<222> (10)...(10)

<400> 38  
Cys Met Leu Gly Arg Val Tyr Arg Pro Cys  
1 5 10

<210> 39  
<211> 11  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> MCH Analog

<221> AMIDATION  
<222> (11)...(11)

<221> DISULFID  
<222> (2)...(11)

<400> 39  
Arg Cys Met Leu Gly Arg Val Tyr Arg Pro Cys  
1 5 10

<210> 40  
<211> 11  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> MCH Analog

<221> MOD\_RES  
<222> (1)...(1)  
<223> Xaa = Des-Amino-Arginine

<221> DISULFID  
<222> (2)...(11)

<221> AMIDATION  
<222> (11)...(11)

<400> 40  
Xaa Cys Met Leu Gly Arg Val Tyr Arg Pro Cys  
1 5 10

<210> 41  
<211> 11  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> MCH Analog

<221> MOD\_RES  
<222> (1)...(1)  
<223> Xaa = D-Arginine

<221> DISULFID  
<222> (2)...(11)

<221> AMIDATION  
<222> (11)...(11)

<221> ACETYLTATION  
<222> (1)...(1)

<400> 41  
Xaa Cys Met Leu Gly Arg Val Tyr Arg Pro Cys  
1 5 10

<210> 42  
<211> 11  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> MCH Analog

<221> MOD\_RES  
<222> (1)...(1)  
<223> Xaa = D-Norleucine

<221> DISULFID  
<222> (2)...(11)

<221> AMIDATION  
<222> (11)...(11)

<221> ACETYLTATION  
<222> (1)...(1)

<400> 42  
Xaa Cys Met Leu Gly Arg Val Tyr Arg Pro Cys  
1 5 10

<210> 43  
<211> 11  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> MCH Analog

<221> MOD\_RES  
<222> (1)...(1)  
<223> Xaa = D-Proline

<221> DISULFID  
<222> (2)...(11)

<221> AMIDATION  
<222> (11)...(11)

<221> ACETYLTATION  
<222> (1)...(1)

<400> 43

Xaa Cys Met Leu Gly Arg Val Tyr Arg Pro Cys  
1 5 10

<210> 44  
<211> 11  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> MCH Analog

<221> MOD\_RES  
<222> (1)...(1)  
<223> Xaa = D-Asparagine

<221> DISULFID  
<222> (2)...(11)

<221> AMIDATION  
<222> (11)...(11)

<221> ACETYLATION  
<222> (1)...(1)

<400> 44  
Xaa Cys Met Leu Gly Arg Val Tyr Arg Pro Cys  
1 5 10

<210> 45  
<211> 11  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> MCH Analog

<221> MOD\_RES  
<222> (1)...(1)  
<223> Xaa = D-Serine

<221> DISULFID  
<222> (2)...(11)

<221> AMIDATION  
<222> (11)...(11)

<221> ACETYLATION  
<222> (1)...(1)

<400> 45  
Xaa Cys Met Leu Gly Arg Val Tyr Arg Pro Cys  
1 5 10

<210> 46  
<211> 11  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> MCH Analog

<221> MOD\_RES

<222> (1)...(1)  
<223> Xaa = D-Glutamic Acid

<221> DISULFID  
<222> (2)...(11)

<221> AMIDATION  
<222> (11)...(11)

<221> ACETYLATION  
<222> (1)...(1)

<400> 46  
Xaa Cys Met Leu Gly Arg Val Tyr Arg Pro Cys  
1 5 10

<210> 47  
<211> 11  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> MCH Analog

<221> AMIDATION  
<222> (11)...(11)

<221> DISULFID  
<222> (2)...(11)

<221> ACETYLATION  
<222> (1)...(1)

<400> 47  
Arg Cys Met Leu Ala Arg Val Tyr Arg Pro Cys  
1 5 10

<210> 48  
<211> 11  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> MCH Analog

<221> ACETYLATION  
<222> (1)...(1)

<221> AMIDATION  
<222> (11)...(11)

<221> DISULFID  
<222> (2)...(11)

<400> 48  
Arg Cys Met Leu Leu Arg Val Tyr Arg Pro Cys  
1 5 10

<210> 49  
<211> 11  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> MCH Analog

<221> ACETYLATION  
<222> (1)...(1)

<221> MOD\_RES  
<222> (5)...(5)  
<223> Xaa = Norleucine

<221> DISULFID  
<222> (2)...(11)

<221> AMIDATION  
<222> (11)...(11)

<400> 49  
Arg Cys Met Leu Xaa Arg Val Tyr Arg Pro Cys  
1 5 10

<210> 50  
<211> 11  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> MCH Analog

<221> ACETYLATION  
<222> (1)...(1)

<221> AMIDATION  
<222> (11)...(11)

<221> DISULFID  
<222> (2)...(11)

<400> 50  
Arg Cys Met Leu Pro Arg Val Tyr Arg Pro Cys  
1 5 10

<210> 51  
<211> 11  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> MCH Analog

<221> ACETYLATION  
<222> (1)...(1)

<221> AMIDATION  
<222> (11)...(11)

<221> DISULFID  
<222> (2)...(11)

<400> 51  
Arg Cys Met Leu Asn Arg Val Tyr Arg Pro Cys  
1 5 10

<210> 52  
<211> 11  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> MCH Analog

<221> ACETYLATION  
<222> (1)...(1)

<221> AMIDATION  
<222> (11)...(11)

<221> DISULFID  
<222> (2)...(11)

<400> 52  
Arg Cys Met Leu Ser Arg Val Tyr Arg Pro Cys  
1 5 10

<210> 53  
<211> 11  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> MCH Analog

<221> ACETYLATION  
<222> (1)...(1)

<221> AMIDATION  
<222> (11)...(11)

<221> DISULFID  
<222> (2)...(11)

<400> 53  
Arg Cys Met Leu Lys Arg Val Tyr Arg Pro Cys  
1 5 10

<210> 54  
<211> 11  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> MCH Analog

<221> ACETYLATION  
<222> (1)...(1)

<221> AMIDATION  
<222> (11)...(11)

<221> DISULFID  
<222> (2)...(11)

<400> 54

Arg Cys Met Leu Glu Arg Val Tyr Arg Pro Cys  
1 5 10

<210> 55  
<211> 11  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> MCH Analog

<221> ACETYLATION  
<222> (1)...(1)

<221> MOD\_RES  
<222> (5)...(5)  
<223> Xaa = D-Leucine

<221> DISULFID  
<222> (2)...(11)

<221> AMIDATION  
<222> (11)...(11)

<400> 55  
Arg Cys Met Leu Xaa Arg Val Tyr Arg Pro Cys  
1 5 10

<210> 56  
<211> 11  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> MCH Analog

<221> ACETYLATION  
<222> (1)...(1)

<221> MOD\_RES  
<222> (5)...(5)  
<223> Xaa = D-Norleucine

<221> DISULFID  
<222> (2)...(11)

<221> AMIDATION  
<222> (11)...(11)

<400> 56  
Arg Cys Met Leu Xaa Arg Val Tyr Arg Pro Cys  
1 5 10

<210> 57  
<211> 11  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> MCH Analog

<221> ACETYLATION

<222> (1)...(1)  
<221> MOD\_RES  
<222> (5)...(5)  
<223> Xaa = D-Proline  
  
<221> DISULFID  
<222> (2)...(11)  
  
<221> AMIDATION  
<222> (11)...(11)  
  
<400> 57  
Arg Cys Met Leu Xaa Arg Val Tyr Arg Pro Cys  
1 5 10  
  
<210> 58  
<211> 11  
<212> PRT  
<213> Artificial Sequence  
  
<220>  
<223> MCH Analog  
  
<221> ACETYLATION  
<222> (1)...(1)  
  
<221> MOD\_RES  
<222> (5)...(5)  
<223> Xaa = D-Glutamic Acid  
  
<221> DISULFID  
<222> (2)...(11)  
  
<221> AMIDATION  
<222> (11)...(11)  
  
<400> 58  
Arg Cys Met Leu Xaa Arg Val Tyr Arg Pro Cys  
1 5 10  
  
<210> 59  
<211> 11  
<212> PRT  
<213> Artificial Sequence  
  
<220>  
<223> MCH Analog  
  
<221> ACETYLATION  
<222> (1)...(1)  
  
<221> MOD\_RES  
<222> (5)...(5)  
<223> Xaa = D-Lysine  
  
<221> DISULFID  
<222> (2)...(11)  
  
<221> AMIDATION  
<222> (11)...(11)

<400> 59  
Arg Cys Met Leu Xaa Arg Val Tyr Arg Pro Cys  
1 5 10

<210> 60  
<211> 11  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> MCH Analog

<221> ACETYLATION  
<222> (1)...(1)

<221> MOD\_RES  
<222> (5)...(5)  
<223> Xaa = bAla

<221> DISULFID  
<222> (2)...(11)

<221> AMIDATION  
<222> (11)...(11)

<400> 60  
Arg Cys Met Leu Xaa Arg Val Tyr Arg Pro Cys  
1 5 10

<210> 61  
<211> 11  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> MCH Analog

<221> ACETYLATION  
<222> (1)...(1)

<221> MOD\_RES  
<222> (5)...(5)  
<223> Xaa = Gamma-Aminobutyric Acid

<221> DISULFID  
<222> (2)...(11)

<221> AMIDATION  
<222> (11)...(11)

<400> 61  
Arg Cys Met Leu Xaa Arg Val Tyr Arg Pro Cys  
1 5 10

<210> 62  
<211> 11  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> MCH Analog

<221> ACETYLATION  
<222> (1)...(1)

<221> AMIDATION  
<222> (11)...(11)

<221> DISULFID  
<222> (2)...(11)

<400> 62  
Arg Cys Met Leu Gly Arg Val Phe Arg Pro Cys  
1 5 10

<210> 63  
<211> 11  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> MCH Analog

<221> ACETYLATION  
<222> (1)...(1)

<221> AMIDATION  
<222> (11)...(11)

<221> DISULFID  
<222> (2)...(11)

<400> 63  
Arg Cys Met Leu Gly Arg Val Trp Arg Pro Cys  
1 5 10

<210> 64  
<211> 11  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> MCH Analog

<221> ACETYLATION  
<222> (1)...(1)

<221> AMIDATION  
<222> (11)...(11)

<221> DISULFID  
<222> (2)...(11)

<400> 64  
Arg Cys Met Leu Gly Arg Val His Arg Pro Cys  
1 5 10

<210> 65  
<211> 11  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> MCH Analog

<221> ACETYLATION  
<222> (1)...(1)  
  
<221> MOD\_RES  
<222> (8)...(8)  
<223> Xaa = (2')Naphthylalanine  
  
<221> DISULFID  
<222> (2)...(11)  
  
<221> AMIDATION  
<222> (11)...(11)  
  
<400> 65  
Arg Cys Met Leu Gly Arg Val Xaa Arg Pro Cys  
1 5 10  
  
<210> 66  
<211> 11  
<212> PRT  
<213> Artificial Sequence  
  
<220>  
<223> MCH Analog  
  
<221> ACETYLATION  
<222> (1)...(1)  
  
<221> MOD\_RES  
<222> (8)...(8)  
<223> Xaa = P-Fluoro-Phenylalanine  
  
<221> DISULFID  
<222> (2)...(11)  
  
<221> AMIDATION  
<222> (11)...(11)  
  
<400> 66  
Arg Cys Met Leu Gly Arg Val Xaa Arg Pro Cys  
1 5 10  
  
<210> 67  
<211> 11  
<212> PRT  
<213> Artificial Sequence  
  
<220>  
<223> MCH Analog  
  
<221> ACETYLATION  
<222> (1)...(1)  
  
<221> MOD\_RES  
<222> (8)...(8)  
<223> Xaa = P-Amino-Phenylalanine  
  
<221> DISULFID  
<222> (2)...(11)  
  
<221> AMIDATION

<222> (11)...(11)

<400> 67

Arg Cys Met Leu Gly Arg Val Xaa Arg Pro Cys  
1 5 10

<210> 68

<211> 11

<212> PRT

<213> Artificial Sequence

<220>

<223> MCH Analog

<221> ACETYLTATION

<222> (1)...(1)

<221> MOD\_RES

<222> (8)...(8)

<223> Xaa = P-Carboxy-Phenylalanine

<221> DISULFID

<222> (2)...(11)

<221> AMIDATION

<222> (11)...(11)

<400> 68

Arg Cys Met Leu Gly Arg Val Xaa Arg Pro Cys  
1 5 10

<210> 69

<211> 11

<212> PRT

<213> Artificial Sequence

<220>

<223> MCH Analog

<221> ACETYLTATION

<222> (1)...(1)

<221> MOD\_RES

<222> (8)...(8)

<223> Xaa - Cyclohexylalanine

<221> DISULFID

<222> (2)...(11)

<221> AMIDATION

<222> (11)...(11)

<400> 69

Arg Cys Met Leu Gly Arg Val Xaa Arg Pro Cys  
1 5 10

<210> 70

<211> 11

<212> PRT

<213> Artificial Sequence

<220>

<223> MCH Analog  
<221> ACETYLATION  
<222> (1)...(1)  
  
<221> MOD\_RES  
<222> (4)...(5)  
<223> Xaa = 5-Aminopentanoic Acid  
  
<221> DISULFID  
<222> (2)...(11)  
  
<221> AMIDATION  
<222> (11)...(11)  
  
<400> 70  
Arg Cys Met Xaa Xaa Arg Val Tyr Arg Pro Cys  
1 5 10  
  
<210> 71  
<211> 11  
<212> PRT  
<213> Artificial Sequence  
  
<220>  
<223> MCH Analog  
  
<221> ACETYLATION  
<222> (1)...(1)  
  
<221> MOD\_RES  
<222> (1)...(1)  
<223> Xaa = D-Arginine  
  
<221> MOD\_RES  
<222> (4)...(5)  
<223> Xaa = 5-Aminopentanoic Acid  
  
<221> DISULFID  
<222> (2)...(11)  
  
<221> AMIDATION  
<222> (11)...(11)  
  
<400> 71  
Xaa Cys Met Xaa Xaa Arg Val Tyr Arg Pro Cys  
1 5 10  
  
<210> 72  
<211> 11  
<212> PRT  
<213> Artificial Sequence  
  
<220>  
<223> MCH Analog  
  
<221> ACETYLATION  
<222> (1)...(1)  
  
<221> MOD\_RES  
<222> (9)...(10)  
<223> Xaa = 5-Aminopentanoic Acid

<221> DISULFID  
<222> (2)...(11)

<221> AMIDATION  
<222> (11)...(11)

<400> 72  
Arg Cys Met Leu Gly Arg Val Tyr Xaa Xaa Cys  
1 5 10

<210> 73  
<211> 11  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> MCH Analog

<221> ACETYLATION  
<222> (1)...(1)

<221> MOD\_RES  
<222> (1)...(1)  
<223> Xaa = D-Arginine

<221> MOD\_RES  
<222> (9)...(10)  
<223> Xaa = 5-Aminopentanoic Acid

<221> DISULFID  
<222> (2)...(11)

<221> AMIDATION  
<222> (11)...(11)

<400> 73  
Xaa Cys Met Leu Gly Arg Val Tyr Xaa Xaa Cys  
1 5 10

<210> 74  
<211> 11  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> MCH Analog

<221> ACETYLATION  
<222> (1)...(1)

<221> MOD\_RES  
<222> (5)...(5)  
<223> Xaa = D-Proline

<221> MOD\_RES  
<222> (9)...(10)  
<223> Xaa = 5-Aminopentanoic Acid

<221> DISULFID  
<222> (2)...(11)

<221> AMIDATION  
<222> (11)...(11)

<400> 74  
Arg Cys Met Leu Xaa Arg Val Tyr Xaa Xaa Cys  
1 5 10

<210> 75  
<211> 11  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> MCH Analog

<221> ACETYLATION  
<222> (1)...(1)

<221> MOD\_RES  
<222> (1)...(1)  
<223> Xaa = Des-Amino-Arginine

<221> MOD\_RES  
<222> (9)...(10)  
<223> Xaa = 5-Aminopentanoic Acid

<221> DISULFID  
<222> (2)...(11)

<221> AMIDATION  
<222> (11)...(11)

<400> 75  
Xaa Cys Met Leu Gly Arg Val Tyr Xaa Xaa Cys  
1 5 10

<210> 76  
<211> 11  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> MCH Analog

<221> ACETYLATION  
<222> (1)...(1)

<221> MOD\_RES  
<222> (4)...(5)  
<223> Xaa = 5-Aminopentanoic Acid

<221> MOD\_RES  
<222> (9)...(10)  
<223> Xaa = 5-Aminopentanoic Acid

<221> DISULFID  
<222> (2)...(11)

<221> AMIDATION  
<222> (11)...(11)

<400> 76

Arg Cys Met Xaa Xaa Arg Val Tyr Xaa Xaa Cys  
1 5 10

<210> 77  
<211> 11  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> MCH Analog

<221> ACETYLATION  
<222> (1)...(1)

<221> MOD\_RES  
<222> (3)...(3)  
<223> Xaa = Norleucine

<221> DISULFID  
<222> (2)...(11)

<221> AMIDATION  
<222> (11)...(11)

<400> 77  
Arg Cys Xaa Leu Gly Arg Val Tyr Arg Pro Cys  
1 5 10

<210> 78  
<211> 11  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> MCH Analog

<221> ACETYLATION  
<222> (1)...(1)

<221> MOD\_RES  
<222> (1)...(1)  
<223> Xaa = Des-Amino-Arginine

<221> MOD\_RES  
<222> (5)...(5)  
<223> Xaa = D-Norleucine

<221> DISULFID  
<222> (2)...(11)

<221> AMIDATION  
<222> (11)...(11)

<400> 78  
Xaa Cys Met Leu Xaa Arg Val Tyr Arg Pro Cys  
1 5 10

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US01/03293

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : C07K 7/08, 7/52; A61K 38/00

US CL : 514/13, 14; 530/317

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
U.S. : 514/13, 14; 530/317, 300

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)  
Please See Continuation Sheet

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DROZDZ et al. Melanin-Concentrating Hormone Binding to Mouse Melanoma Cells in Vitro. FEBS Letters. 1995, Vol. 359, pages 199-202. See pages 200-202, Figure 1	1, 9-11
X	DROZDZ et al. (D-(p-Benzoylphenylalanine)13Tyrosin19)-Melanin-Concentrating Hormone, a Potent Analogue for MCH Receptor Crosslinking. J. Peptide Science. 1999, Vol. 5, pages 234-242. See pages 236-239, Figure 1.	1, 9-11
X	HINTERMANN et al. Synthesis and Characterization of New Radioligands for the Mammalian Melanin-Concentrating Hormone (MCH) Receptor. J. Receptor & Signal Transduction Research. 1999, Vol. 19, pages 411-422. See pages 414-418, Figures 1-3.	1, 9-11
X	BAKER et al. Structure-Activity Studies with Fragments and Analogues of Salmonid Melanin-Concentrating Hormone. Peptides (Fayetteville, N. Y.) 1990, Vol. 16, pages 1103-1108. See pages 1104-1106, Table 1.	1
X	LEBL et al. Melanin Concentrating Hormone Analogues: Contraction of the Cyclic Structure.1. Agonist Activity. J. Med. Chem. 1988, Vol. 31, pages 949-954. See Figures 1 and 2.	1
X	WO 90/11295 A1 (THE SALK INSTITUTE FOR BIOLOGICAL STUDIES) 04 October 1990 (04,10,1990). See pages 3, 6 and 16, Example 1.	1

☐ Further documents are listed in the continuation of Box C.

☐ See patent family annex.

Special categories of cited documents:	
"A" document defining the general state of the art which is not considered to be of particular relevance	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"E" earlier application or patent published on or after the international filing date	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"O" document referring to an oral disclosure, use, exhibition or other means	"Z" document member of the same patent family
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

04 May 2001 (04.05.2001)

Date of mailing of the international search report

24 MAY 2001

Name and mailing address of the ISA/US

Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703)305-3230

Authorized officer

Chih-Min Kuan

Telephone No. (703) 308-0196

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US01/03293

## Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claim Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claim Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claim Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:  
Please see Continuation Sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.: 1-12, SEQ ID NO:8

Remark on Protest

☐  
☐

The additional search fees were accompanied by the applicant's protest.

No protest accompanied the payment of additional search fees.

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US01/03293

**Continuation of B. FIELDS SEARCHED Item 3:** STN search on CAPLUS, MEDLINE, EMBASE, BIOSIS and SCISEARCH; EAST search on USPAT, DERWENT, EPO, JPO. Search term used: melanin concentrating hormone, MCH, melanin concentrating hormone receptor, MCH receptor, binding, inhibiting, agonist antagonist. Amino acid sequence search on SEQ ID NO: 7, 8, 9, 10.

### BOX II Observations where unity of invention is lacking

This International Searching Authority found multiple inventions in this international application, as follows:

This International Search Authority has found 176 inventions claimed in the International Application covered by the claims indicated below:

Group 1-44, claims 1-8 and 9-12, in part drawn to one of peptides SEQ ID NO: 7, 8, 9, 10, 15, 24, 25, 27, 28, 30-49, 51, 52, 56, 61, 62, 63, 65-67, 69-72 and 77, respectively and a method of screening for a compound able to bind a MCH receptor using such. For group 1, the claims will be searched to extent that they read upon SEQ ID NO: 7. If group 2 is elected, the claims will be searched to extent that they read upon SEQ ID NO: 8.

Group 45-88, claim 13, in part drawn to a method for increasing weight in a subject comprising the step of administering one of peptides SEQ ID NO: 7, 8, 9, 10, 15, 24, 25, 27, 28, 30-49, 51, 52, 56, 61, 62, 63, 65-67, 69-72 and 77, respectively. For group 45, the claims will be searched to extent that they read upon SEQ ID NO: 7. If group 46 is elected, the claims will be searched to extent that they read upon SEQ ID NO: 8.

Group 89-132, claim 14, in part drawn to a method for increasing appetite in a subject comprising the step of administering one of peptides SEQ ID NO: 7, 8, 9, 10, 15, 24, 25, 27, 28, 30-49, 51, 52, 56, 61, 62, 63, 65-67, 69-72 and 77, respectively. For group 89, the claims will be searched to extent that they read upon SEQ ID NO: 7. If group 90 is elected, the claims will be searched to extent that they read upon SEQ ID NO: 8.

Group 133-176, claims 15, and 16, in part drawn to a method for measuring the ability of a compound to decrease weight or appetite in a subject comprising the step of administering one of peptides SEQ ID NO: 7, 8, 9, 10, 15, 24, 25, 27, 28, 30-49, 51, 52, 56, 61, 62, 63, 65-67, 69-72 and 77, respectively. For group 133, the claims will be searched to extent that they read upon SEQ ID NO: 7. If group 134 is elected, the claims will be searched to extent that they read upon SEQ ID NO: 8.

The International Search Authority considers that the international application does not comply with the requirements of unity of invention (Rules 13.1, 13.2 and 13.3) for the reasons indicated below:

The inventions listed as Groups 1-176 do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons:

The peptides of SEQ ID NO: 7, 8, 9, 10, 15, 24, 25, 27, 28, 30-49, 51, 52, 56, 61, 62, 63, 65-67, 69-72 and 77 are unrelated, each to the other. Each peptide is distinct and does not share a special technical feature with the other. Additionally, the claimed methods produce different results which are not coextensive and which do not share the same technical feature.